

- littoralis.** 22. *M. nigra*, thorace rufo, abdomine anteriori macula biloba alba: segmentorum quinque posteriorum marginibus albis. *Petagn. itif. Cal. p. 33. n. 171. f. 37.*
Habitat in littore Brancalioni Calabriae, europaeae affinis, brutta minor, maxillis porrectis oculisque nitidis, antennis longitudine capitis pedibusque testaceis.
- spinosa.** 23. *M. flavescens nigro varia*, thorace utrinque spinis quatuor, abdominis segmento secundo maculis duabus testaceis. *Swederus nov. act. Stockh. 8. 1787. 4. n. 2. 39.*
Habitat in Rio Janeiro, hirsuta, at inter maximas hujus generis, americanae facie.
- cephalotes.** 24. *M. capite magno aterrimo utrinque spinoso*, abdomine macula aurea fasciisque tribus interruptis albis. *Swederus nov. act. Stockh. 8. 1787. 4. n. 2. 40.*
Habitat in Georgia Americae, valde pubescens, europaea triplo fere major.
- bifasciata.** 25. *M. atra*, capite et thorace supra, abdominisque fasciis duabus rubris, alis violaceo-nigris. *Swederus nov. act. Stockh. 8. 1787. 4. n. 2. 41. 42.*
Habitat in Noveboraco, tota pubescens, europaea duplo major.
- sexpunctata.** 26. *M. nigra*, thorace subcylindrico immaculato: abdomine punctis utrinque tribus albidis. *Swederus nov. act. Stockh. 8. 1787. 4. n. 2. 43.*
Habitat in Africa, aptera, albo-subpubescens, europaea triplo fere minor.
- sexmaculata.** 27. *M. nigra*, thorace supra rufo, abdomine utrinque serie macularum trium albidarum. *Swederus nov. act. Stockh. 8. 1787. 4. n. 2. 44.*
Habitat in India, aptera, pubescens, europaea duplo fere major.