

most southern point in which the shining slave maker occurs in the Appalachians.

86. *Camponotus castaneus* Latr. Clemson College (M. R. S.); Florence (O. L. C.); Pinewood (D. E. R.).
87. *Camponotus castaneus* subsp. *americanus* Mayr. Clemson College, Walhalla (M. R. S.); Abbeville, Spartanburg (D. E. R.).
88. *Camponotus herculeanus* subsp. *pennsylvanicus* DeGeer. Clemson College (M. R. S.); Walhalla (J. E. W.); Rocky Bottom (W. C. N.); Florence (O. L. C.); Pelzer, Conway (D. E. R.). This is the common black carpenter ant, which is fond of nesting in logs, stumps, trees, etc.
89. *Camponotus caryæ* Fitch. Clemson College (J. E. W.); Westminster, Spartanburg, Batesburg, Camden, Ridgeland, Conway, McCall (D. E. R.).
90. *Camponotus caryæ* subsp. *rasilis* Whlr. Pawleys Island (O. L. C.); Columbia (D. E. R.).
91. *Camponotus caryæ rasilis* var. *decipiens* Emery. Clemson College (F. M.); Pickens, Anderson, Laurens, Spartanburg, Inman, Cowpens, Union, Edgefield, Johnson (D. E. R.).
92. *Camponotus caryæ rasilis* var. *pavidus* Whlr. Due West (D. E. R.).
93. *Camponotus abdominalis* subsp. *floridanus* Buckley. Hilton Head Island (H. L. S.). This ant, perhaps the commonest species of *Camponotus* in Florida, certainly ranges as far north as North Carolina.
94. *Camponotus (Colobopsis) mississippiensis* M. R. Smith. Adams Run (D. E. R.). In Mississippi this species generally nests in the twigs of white ash.
95. *Camponotus (Colobopsis) pylartes* subsp. *fraxinicola* M. R. Smith. Charleston (D. L.); Williamston (D. E. R.)
96. *Camponotus (Colobopsis) obliquus* M. R. Smith. Spartanburg (D. E. R.).