

2. *Myrmica rubra* L. subsp. *brevinodis* n. subsp. var. *sulcinodoides* n. var. (*Myrmica brevinodis* Emery)
 ♀. S. Dakota. One of the original localities. 8: 313 (1895). For type locality of *brevinodis*, see above.
3. *Myrmica rubra* L. subsp. *scabrinodis* Nyl. var. *fracticornis* n. var. (*Myrmica lobicornis fracticornis* Emery)
 ♀ ♀. Dakota. 8: 313-314 (1895). According to Emery the Dakota specimens represent a transitory form between the var. *fracticornis* and the var. *sabuleti*.
 Type locality of *fracticornis*, Buffalo, N. Y.
4. *Myrmica rubra* L. subsp. *scabrinodis* Nyl. var. *sabuleti* Meinert (*Myrmica americana* Weber)
 Presumably ♀ ♀ ♂. S. Dakota. One of several localities. 8: 314 (1895). Emery remarked that other workers from S. Dakota are smaller and darker colored than *sabuleti* and are therefore much nearer the var. *schrencki*.
 Type locality of *americana*, Colebrook, Conn.
5. *Crematogaster lineolata* Say subsp. *lineolata* Say typical var. *cerasi* Fitch (*Crematogaster (Acrocoelia) lineolata* (Say))
 ♀. Dakota. One of several localities. 8: 282-283 (1895).
 Type locality of *cerasi*, N. Y., that of *lineolata*, Ind.
6. *Solenopsis molesta* Say (*Solenopsis (Diplorhoptrum) molesta* (Say))
 ♀ ♀ ♂. Dakota. One of several localities. 8: 277-278 (1895). According to Emery females from S. Dakota are slightly darker than the typical form.
 Type locality of *molesta*, Ind.
7. *Leptothorax (Leptothorax) muscorum* Nyl. var. (*Leptothorax (Mychothorax) canadensis yankee* Emery)
 1 ♀. Hill City, S. Dakota. 8: 317-318 (1895). Wheeler considered the above-mentioned specimen to belong to his *muscorum* var. *sordidus*, the type locality of which was Boulder, Colo.
 For type locality of *yankee*, see below.
- *8. *Leptothorax (Leptothorax) canadensis* Prov. var. *yankee* n. var. (*Leptothorax (Mychothorax) canadensis yankee* Emery)