

14. *Camponotus herculeanus* L.
No castes mentioned by Emery. Dakota. One of several localities. 7: 674 (1893).
Apparently the form or forms Emery had before him were what Creighton (1950) calls *pennsylvanicus* (Degeer) and *pennsylvanicus modoc* Wheeler. Type locality of *pennsylvanicus*, Pa., of *modoc*, Calif.
15. *Lasius (Lasius) niger* L. var. *neoniger* n. var. (*Lasius (Lasius) niger neoniger* Emery)
♂. Hill City, S. Dakota. One of several localities. 7: 639 (1893). Emery remarks that only a few workers from Hill City agree fairly well in color and pilosity with the typical *niger* but are smaller than the common Palearctic form.
Type locality for *neoniger* has not been designated.
16. *Lasius (Lasius) brevicornis* n. sp. (*Lasius (Chthonolasius) brevicornis* Emery)
♂ ♀ ♂. Dakota. One of the original localities. 7: 637-640 (1893).
Type locality of *brevicornis*, D. C.
17. *Lasius (Lasius) flavus* L. (*Lasius (Chthonolasius) flavus nearcticus* Wheeler)
♂ ♀. Dakota. One of several localities. 7: 637-638 (1893).
Type locality of *nearcticus*, Ill.
18. *Lasius (Acanthomyops) claviger* Roger (*Acanthomyops claviger* (Roger))
♂ ♀. Dakota. One of several localities. 7: 638, 642 (1893).
Type locality of *claviger*, Pa.
19. *Lasius (Acanthomyops) interjectus* Mayr (*Acanthomyops interjectus* (Mayr))
No caste or castes indicated. Dakota. One of several localities. 7: 638, 642 (1893).
Type locality of *interjectus*, N. J.
20. *Formica (Formica) fusca* L. subsp. *subpolita* Mayr var. *neogagates* n. var. (*Formica (Proformica) neogagates* Emery)