

- Presumably ♀ ♀ ♂. Castes not definitely indicated. Dakota. One of the original localities. 7: 646, 661-662 (1893). Emery mentions workers and a female from Hill City that are not entirely typical.
Type locality of *neogagates*, Beatty, Pa.
- *21. *Formica (Formica) lasioides* n. sp. (*Formica (Proformica) lasioides* Emery)
3 ♀. Hill City, S. Dakota, from Mr. Pergande. Only locality mentioned. 7: 646, 664 (1893).
Type locality of *lasioides*, Hill City, S. D.
22. *Formica lasioides* Em. var. *picea* n. var. (*Formica (Proformica) lasioides* Emery)
♀. Hill City, S. Dakota, from Mr. Pergande. One of two localities mentioned. 8: 335 (1895). According to Emery the workers from Hill City form a transition from *picea* to *lasioides*.
Type locality of *vetula* (n. name for *picea*), Yale, B. C., for *lasioides*, see above.
23. *Formica (Formica) pallide-fulva* Latr. subsp. *fusca* n. subsp. (*Formica (Neoformica) pallide-fulva nitidiventris* Emery)
♀ ♀. Hill City, S. Dakota. One of two localities mentioned. 7: 645, 656 (1893). According to Emery somewhat lighter and shining specimens, including a female from Hill City form a transition to *nitidiventris*.
Type locality of *nitidiventris*, D. C.
- *24. *Formica (Formica) fusca* L. var. *subaenescens* n. var. (*Formica (Formica) fusca* L.)
♀. S. Dakota. One of the original localities. 7: 646, 659-660 (1893). Type locality of *subaenescens*, S. D., of *fusca*, Europe.
- *25. *Formica (Formica) fusca* L. var. *neorufibarbis* n. var. (*Formica (Formica) neorufibarbis* Emery)
♀. S. Dakota. One of the original localities. 7: 646, 660 (1893).
Type locality of *neorufibarbis*, Hill City, S. D.
- *26. *Formica (Formica) dakotensis* n. sp. (*Formica (Formica) dakotensis* Emery)