

- ♂. Hill City, S. Dakota, from Mr. Pergande. Only locality mentioned by Emery. 7: 644, 652-653 (1893).
Type locality of *dakotensis*, Hill City, S. D.
27. *Formica (Formica) fusca* L. *subpolita* Mayr var.? *specularis* n. var. (*Formica (Formica) dakotensis* Emery)
♀. Dakota, received from Mr. Pergande. One of the original localities. 7: 663 (1893).
Type locality of *specularis*, Wis., of *dakotensis*, see above.
28. *Formica (Formica) rufa* L. subsp. *integra* Nyl. var. *hæmorrhoidalis* n. var. (*Formica (Formica) integra hæmorrhoidalis* Emery)
♂. Dakota. One of the two original localities. 7: 644, 652 (1893).
Type locality of *hæmorrhoidalis*, Colo.
29. *Formica (Formica) rufa* L. subsp. *obscuriventris* Mayr var. *rubiginosa* n. var. (*Formica (Formica) obscuripes* Forel)
♂. Dakota. One of the original localities. 7: 644, 650 (1893). Creighton, 1940, (Amer. Mus. Novitates No. 1055, p. 5) suggested that Emery had a mixed series when describing *rubiginosa*: *Formica rufa obscuripes* Forel (Colo.), *F. rufa integra* Nyl. (Nebr.), *F. rufa melanotica* Emery and *F. rufa gymnomma* Wheeler (S. Dak.). Creighton (1950) considers both *rubiginosa* and *melanotica* a synonym of *obscuripes*, and *gymnomma* a synonym of *obscuriventris* Mayr. The type locality for *obscuripes*, Green River, Wyo., for *melanotica*, Wis., for *gymnomma*, Cold Spring Harbor, Long Island, N. Y., and for *obscuriventris*, Conn.
- *30. *Formica (Formica) ulkei* n. sp. (*Formica (Formica) ulkei* Emery)
♂. Hill City, S. Dakota, from Mr. Pergande. Only locality mentioned. 7: 643, 653-654 (1893).
Type locality of *ulkei*, Hill City, S. D.
- *31. *Formica (Formica) sanguinea* Latr. subsp. *puberula* n. subsp. (*Formica (Raptiformica) puberula* Emery)
1 ♂. Hill City, S. Dakota. 7: 643, 648 (1893). Other specimens from Colorado Emery considered as a transitory