

- Head subopaque. Middle and hind tibia each without a row of graduated, erect, bristles on their flexor surface (fig. 34).  
*Camponotus tortuganus* Emery (p. 62).
9. Eye located nearer to the base of the mandible than to the posterior border of the head----- 10  
Eye located nearer to the posterior border of the head than to the base of the mandible----- 11
10. Antennae and legs extraordinarily long. Body slender and bearing long coarse, suberect to erect, grayish or whitish hairs; the suberect to erect hairs normally absent from the scapes. Integument with a peculiar grayish-violaceous luster or sheen (fig. 41). Introduced species.  
*Paratrechina longicornis* (Latreille) (p. 74).  
Unlike alternative above in one or more characters. Scapes usually, and tibiae always, with coarse, suberect to erect hairs (fig. 42)----*Paratrechina* (Nylanderia) spp. (p. 76).
11. Thorax with a strong constriction in the mesonotum, the constriction somewhat subcylindrical. Scape extending approximately one-half its length beyond the posterior border of the head. Gaster from above, with the base meeting each side in a distinct angle (fig. 43).  
*Prenolepis imparis* (Say) (p. 78).  
Differing from the above in one or more characters----- 12
12. Maxillary palpus short, 3-segmented. Eye unusually small to small. Body hairs commonly barbed.<sup>1</sup> Workers capable of emitting pleasant lemon-verbena or citronella odors-- 13  
Maxillary palpus long, 6-segmented. Eye normal. Body hairs not barbed. Workers capable of emitting distinct formic acid odors----- 16
13. Petiolar node, in profile, with a narrow and rather sharp summit (fig. 48,a)----- 14  
Petiolar node, in profile, with a thicker, blunt summit (fig. 49)----- 15
14. Antennal scape when fully extended with its apex considerably surpassing the posterior border of the head. A small tooth on the superior border of the mandible near the junction of the mandible's superior and masticatory border (fig. 48, b). Erect hairs on the dorsal surface of the gaster largely confined to a transverse row on the posterior border of each segment (fig. 48, a).  
*Acanthomyops interjectus* (Mayr) (p. 88).  
Antennal scape when fully extended with its apex either not attaining or else not surpassing the posterior border of the head. The small tooth lacking on the superior border of the mandible as described for *interjectus*. Dorsal surface of gaster with widely distributed, abundant, erect hairs (fig. 47)-----*Acanthomyops claviger* (Roger) (p. 86).

<sup>1</sup> It may be difficult to see the barbs on the hairs except under high magnification: not less than 61.2.