

2. *Odontomachus haematoda* (L.)

Tropicopolitan; large, dark brown and stinging painfully; generally carnivorous.

3. *Pheidole*, near *rinae tipuna* Forel

Although Wheeler recorded *P. javana* Mayr from Guam, the present species in the soldier caste is much smaller and less glabrous. It is also much smaller than *oceanica* and *bolabolensis*. The Saipan soldier is close to *rinae* subsp. *tipuna* Forel but has a distinctly shorter head and is also close to the *rinae* subsp. *incensa* Wheeler types in the American Museum of Natural History.

4. *Pheidole* sp.

Workers of a second species, impossible of identification without the soldier caste, differ in being smooth, shiny and dark rather than being densely punctate and pale.

5. *Cardiocondyla emeryi* Forel

One dealate female of this cosmopolitan species.

6. *Vollenhovia pedestris* F. Smith

Not in the Guam or Bikini lists. The workers agree well with Mann's British Solomon specimens; subspecies are known from New Guinea, the Seychelles, Borneo, the Philippines and other paleotropical localities.

7. *Monomorium destructor* (Jerdon)

Tropicopolitan; tiny and reddish yellow.

8. *Monomorium floricola* (Jerdon)

Tropicopolitan; even smaller than the preceding and a shiny dark brown in color.

9. *Solenopsis geminata* subspecies *rufa* (Jerdon)

Showing the mesosternal spine characteristic of this widespread paleotropical form of the cosmopolitan species. I have