

of *Lioponera*, *Sphinctomyrmex*, etc., like the normal winged females of the majority of the Ponerinæ, lead very naturally to the conditions seen in the more modified winged females of the Myrmicinæ, Dolichoderinæ and Camponotinæ.

## BIBLIOGRAPHY.

**Emery, C.**

- '95 Die Gattung *Dorylus* Fab. und die systematische Eintheilung der Formiciden. Zool. Jahrb. (Abth. f. Syst.), 8 Bd., 1895, pp. 686-778. Taf. 14-17, 41 text-figs.
- '01a Notes sur les Sous-familles des Dorylines et Ponerines (Famille des Formicides). Ann. Soc. Ent. Belg., Tome XLV., 1901, pp. 32-54.
- '01b Note mirmecologique. I. Revisione del Gruppo dei Generi Affini a *Cerapachys* F. Sm. Rend. Sess.R. Accad. Sci. Ist. Bologna, Anno 1901, 8 pp.

**Forel, A.**

- '99 Formicidæ (in *Biologia Centrali-Americana*), 1899, pp. 2 and 22.
- '01 À propos de la Classification des Fourmis. Ann. Soc. Ent. Belg., Tome XLV., 1901, pp. 136-141.

**Wheeler, W. M.**

- '00 The Habits of *Ponera* and *Stigmatomma*. Biol. Bull., Vol. II., No. 2, Nov., 1900.

ROCKFORD, ILLINOIS, July 6, 1902.