

33. *Strumigenys clypeata* Roger. "Carolina." (Roger.)

34. *Atta* (*Trachymyrmex*) *septentrionalis* McCook [= *tardigrada* (Buckl.) auct.]. Black Mountain. (Forel.)

SUBFAMILY DOLICHODERINÆ.

35. *Dolichoderus plagiatus* Mayr var. *beutenmülleri* var. nov. Differs from the typical form in the total absence of spots on the basal gastric segments. Eleven workers taken by beating, near the Black Mountain R.R. Station.

36. *Dolichoderus mariæ* Forel. A large colony, comprising several hundred workers of this beautiful species, found under a small stone in the pine woods at Panther Gap, Blue Ridge, Buncombe Co.

37. *Dolichoderus taschenbergi* Mayr. Emery mentions specimens from "Carolina" in the Berlin Museum.

38. *Dorymyrmex pyramicus* Roger var. *niger* Pergande. Faisons, N. G. (Forel.)

39. *Dorymyrmex pyramicus* Roger var. *flavus* McCook. Faisons, N. C. (Forel.) Forel describes a mixed colony of this and the preceding variety.

40. *Tapinoma sessile* (Say) Emery. Workers from four colonies of the typical dark-colored form of this species. Different localities in the Swannanoa Valley, Black Mountain, etc.

41. *Iridomyrmex analis* André. This is, in all probability, the form mentioned by Forel as occurring near Morganton, N. C., and as being related to *I. humilis*.

SUBFAMILY CAMPONOTINÆ.

42. *Brachymyrmex heeri depilis* Emery. Several workers from the North Fork of the Swannanoa River.

43. *Prenolepis imparis* (Say) Emery var. *testacea* Emery. Numerous workers from six different colonies of this variety,