

- Brownish black... (41) *fallax subbarbatus* var. *paucipilis* Emery.
52. Head blackish brown..... 53
 Head and thorax yellowish red..... (42) *fallax discolor* Emery.
53. Thorax blackish.... (44) *fallax discolor* var. *cnemidatus* Emery.
 Thorax red..... (43) *fallax discolor* var. *clarithorax* Emery.
54. Thorax feebly and evenly arcuate above, without a distinct impression at the mesoëpinal suture; border of truncated surface of head sharp.... (54) *abditus* var. *etiolatus* Wheeler.
 Thorax, with a distinct impression at the mesoëpinal suture; border of truncated surface of head blunt..... 55
55. Gaster entirely black..... (55) *impressus* Roger.
 Gaster banded with yellow at the base..... 56
56. Thorax and posterior portion of head dark brown.. (56) *pylartes* Wheeler.
 Head and thorax yellow..... (57) *pylartes* var. *hunteri* var. nov.

A. SUBGENUS CAMPONOTUS

I. *Maculatus* Group1. *Camponotus maculatus vicinus* Mayr

Camponotus vicinus MAYR, Verh. zool. bot. Ges. Wien, XX, p. 940, ♀, 1870; FOREL, Bull. Soc. Vaud. Sci. Nat., (2) XVI, p. 60, ♀, 1879; DALLA TORRE, Catalog. Hymen., VII, p. 257, 1893.

C. sylvaticus var. *vicinus* MAYR, Verh. zool. bot. Ges. Wien, XXXVI, p. 422, 1886.

C. maculatus subsp. *vicinus* EMERY, Zool. Jahrb. Abth. f. Syst., VII, p. 671, ♀ ♀, 1893; FOREL, Biol. Centr. Amer., p. 131, 1899-1900.

Worker major. Length, 11-13 mm.; head, 3.7 x 3.5 mm.; scape, 3.2 mm.; hind tibia, 3.3 mm.

Head, excluding the mandibles, nearly as broad as long, broader behind than in front, with feebly concave posterior and feebly convex lateral borders, convex dorsal and flattened gular surfaces. Mandibles strong, convex, 6-7-toothed. Clypeus carinate, its anterior border moderately produced in the form of a broad flap-like lobe with rounded corners and crenate edge. Frontal carinae lyrate. Eyes moderately large, flattened. Antennal scapes distinctly flattened at the base and but slightly widened at their tips, which reach a short distance beyond the posterior corners of the head. Thorax robust in front though narrower than the head; pro- and mesonotum slightly flattened; meso- and metapleural regions compressed; in profile, the dorsum is evenly arched, the epinotum with subequal base and declivity, forming at their juncture a rounded, obtuse angle. Petiole with convex anterior and flattened posterior surface and rather blunt border. Gaster of the usual shape. Legs long; middle and hind tibiae elliptical in cross-section, neither compressed nor grooved.

Mandibles and anterior borders of cheeks shining, the former very coarsely striato-punctate; head, thorax and gaster subopaque; head densely shagreened