

**IV. *Fallax Group***

As I have recently published the synonymy and descriptions of the twelve North American subspecies and varieties of *C. fallax* Nyl., which is the type of this group (Journ. N. Y. Ent. Soc., XVIII, No. 4, 1910), I shall here confine myself to listing the various forms, with the states in which they are known to occur.

**33. *Camponotus fallax* Nyl. var. *nearcticus* Emery**

New York; New Jersey; Pennsylvania; Connecticut; Massachusetts; Rhode Island; Illinois; Wisconsin; Nebraska; Washington; Idaho; Oregon; California; Florida; Texas; Canada.

**34. *C. fallax* var. *minutus* Emery**

New Jersey; New York; Pennsylvania; Massachusetts; Illinois; Canada; Vancouver.

**35. *C. fallax* var. *pardus* Wheeler**

New York; New Jersey.

**36. *C. fallax* var. *tanquaryi* Wheeler**

Illinois.

**37. *C. fallax* var. *decipiens* Emery**

Indiana; Kansas; Colorado; Utah.

**38. *C. fallax rasilis* Wheeler**

Texas; Arizona; Louisiana; Florida.

**39. *C. fallax rasilis* var. *pavidus* Wheeler**

Texas; Louisiana; Florida.

**40. *C. fallax subbarbatus* Emery**

Maryland; Virginia; New Jersey; Illinois; California.

**41. *C. fallax subbarbatus* var. *paucipilis* Emery**

Maryland.

**42. *C. fallax discolor* Buckley**

Texas; Oklahoma; Missouri; Illinois.