

Subfamily MYRMICINÆ.

8. *Cardiocondyla emeryi* Forel.

A single worker of this species, which has not before been recorded from the United States, was sent me by the American Museum of Natural History from Miami, Florida. The species is widely distributed in the West Indies (St. Thomas, St. Vincent, Bahamas, Cuba, Porto Rico, Jamaica) and in several localities in the Old World tropics (Palestine, India, Madagascar, Polynesia).

9. *Solenopsis picta* Emery var. *moerens* var. nov.

Worker. Differing from the type of the species from Florida only in color, the whole body being uniform piceous brown, the mandibles, antennæ and legs brownish yellow.

Six workers taken by Mr. J. D. Mitchell from a dead pecan twig at Victoria, Texas.

10. *Solenopsis huachucana* sp. nov.

Worker. Length 1.5-2 mm.

Head subrectangular, distinctly longer than broad, a little broader in front than behind, with rounded posterior corners and nearly straight posterior and lateral borders. Eyes large, convex, with 6-8 facets in their greatest longitudinal diameter. Clypeus prominent, with strong median carinæ terminating at the anterior margin in two large, acute teeth; lateral teeth small and blunt, indistinct in some specimens. Mandibles with oblique 4-toothed apical margins. Antennæ slender, scapes reaching $\frac{2}{3}$ the distance from the eyes to the posterior corners of the head, club not longer than the remainder of the funiculus, but much broader, its basal fully half as long as its terminal joint; first and second joints much longer than broad, joints 3-8 fully as long as broad, at least in large individuals. Thorax through the pronotum a little more than half as broad as the head; in profile the dorsal outline of the pronotum is convex in front and straight and sloping behind to the short but pronounced mesoëpinal constriction; epinotum with the base longer than the declivity, the former convex in profile, the latter sloping and concave, the angle between the two surfaces obtuse and much rounded. Petiole with a very short, narrow peduncle, the node slightly broader than long; postpetiole slightly lower and broader, nearly spherical above, a little broader than long; neither petiole nor postpetiole with a distinct tooth on the ventral side. Gaster elliptical, its anterior border concave in the middle.

Smooth and shining throughout, with very sparse and indistinct piligerous punctures on the body and legs.

Hairs whitish, bristly, of unequal length, erect on the body and moderately abundant, shorter on the antennal scapes and legs.