

Kauai: Lihue (type locality). Oahu (Van Zwaluwenburg): Honolulu (Bryan); Moanalua; Palolo; Tantalus; Waianae (Illingworth); Hauula; Kolekole Pass. Hawaii: Hilo Sugar Company (Swezey). Maui: Hana (Bryan). Molokai (Illingworth).

French Frigates Shoal (Bryan) ♀. Necker Island, ♂. Laysan Island (Fullaway) ♀.

4. *Ponera perkinsi* Forel.

Ponera perkinsi Forel: Fauna Hawaiiensis, vol. 1, p. 117, 1899, ♀ ♀ ♂. Wheeler: Lab. Zool. Gen. Agrar. Portici, Boll., vol. 3, p. 271, 1909, ergatomorphic ♂. Haw. Ent. Soc., Proc.: Gulick, vol. 2, pp. 306, 310, 1913, appendix; Perkins, vol. 1, p. 43, 1906; Illingworth, vol. 5, p. 273, 1923; Swezey and Bryan, vol. 7, p. 296, 1929.

Oahu: Tantalus, Lanihuli (Swezey, Bryan); Honolulu (Bridwell, Illingworth); Kuliouou, Palolo Valley, Kaumuahona, Kaimuki, Niu (Swezey); Nuuanu Pali (Ehrhorn). Molokai: forests (Swezey and Bryan). Maui: Haleakala (Swezey). Kauai: Summit Camp (Swezey); Koloa. Hawaii: Waimea.

"In the mountains on all the islands, generally from 2,000 to 4,000 feet" (Perkins).

5. *Ponera punctatissima schauinslandi* Emery.

Ponera punctatissima schauinslandi Emery: Zool. Jahrb., Syst., vol. 12, p. 439, 1899, ♀.

Laysan Island: type locality (Schauinsland).

6. *Pseudocryptopone zwaluwenburgi* Wheeler.

Pseudocryptopone zwaluwenburgi Wheeler: Am. Mus., Novitates, no. 672, p. 14, fig. 5, 1933, ♀.

Oahu: Honolulu (type locality); Waimanalo, Waialua, Waipio (Van Zwaluwenburg), in soil of sugar cane fields, both fallow and under cultivation. Maui: central portion, in field of growing sugar cane (Van Zwaluwenburg).

7. *Pseudocryptopone swezeyi* Wheeler.

Pseudocryptopone swezeyi Wheeler: Am. Mus., Novitates, no. 672, p. 16, fig. 6, 1933, ♀ ♀.

Oahu: Honolulu (type locality), in soil of sugar cane fields, both fallow and under cultivation (Van Zwaluwenburg).