



FIG. 8-11.—Scanning electron micrographs of mechanoreceptors and chemoreceptors of the venom apparatus and adjacent 7th sternum and remnant of 10th abdominal segment. 8. Hair plate on the oblong plate (OP) at its point of articulation with the triangular plate (1000 \times). 9. Seventh sternum (7 ST), showing the numerous trichoid sensilla and laterally arranged chemosensitive basiciconic sensory spine (SS) (100 \times). 10. Enlargement of one sensory spine on 7th sternum, showing the pore at its distal tip (1000 \times). 11. Anal pad (AP), showing the long trichoid sensilla extending down past the anal area toward the lower venom structures (100 \times). Note: Figures reduced to 90%.

REFERENCES CITED

- Dethier, V. G. 1957. Chemoreception and the behavior of insects. *Surv. Biol. Prog.* 3: 149-83.
- Hermann, H. R. 1968a. The hymenopterous poison apparatus. IV. *Dasymutilla occidentalis* (Hymenoptera: Mutillidae). *J. Georgia Entomol. Soc.* 3: 1-10.
- 1968b. The hymenopterous poison apparatus. V. *Ancurectus simoni*. *Ann. Entomol. Soc. Amer.* 61: 1315-7.
1969. The hymenopterous poison apparatus: Evolutionary trends in three closely related subfamilies of ants (Hymenoptera: Formicidae). *J. Georgia Entomol. Soc.* 4: 123-41.
1976. The furcula, a major component of the hymenopterous venom apparatus. *Int. J. Insect Morphol. Embryol.*
- Hermann, H. R., and M. S. Blum. 1966. The morphology and histology of the hymenopterous poison apparatus. I. *Paraponera clavata* (Formicidae). *Ann. Entomol. Soc. Am.* 59: 397-409.