

BOCAGIANA

Museu Municipal do Funchal

Madeira

1.VII.1993

No. 167

***MYRMECINA GRAMINICOLA* (LATR., 1802)
(HYMENOPTERA, FORMICIDAE) IN MADEIRA**

By X. ESPADALER* & M. BÁEZ**

SUMMARY. Two males of *Myrmecina graminicola* (LATR., 1802) were found at Portela, Madeira. This ant is to be added to the fifteen ant taxa known from Madeira, that is the southernmost locality for this species.

SUMÁRIO. Dois machos de *Myrmecina graminicola* foram encontrados na Portela, Madeira. Esta espécie de formiga é acrescentada aos quinze taxa de formigas conhecidas para a Madeira, que é a localidade mais meridional para esta espécie.

Only fifteen ant taxa are known to occur on the archipelago of Madeira (WHEELER, 1927; WELLENIUS, 1949); the small number of ant species on oceanic islands is a well known phenomenon (TAYLOR & WILSON, 1961; WILSON, 1973; HEINZE, 1986); any addition to the checklist is worth mention.

During a short visit to the Island, one of the authors (M.B.) got two males of a genus never reported for the archipelago: *Myrmecina graminicola* (LATREILLE, 1802). Both males were captured by sweeping vegetation at 16 hrs at a partially cultivated field near a laurel forest at Portela, Madeira, the 27 August, 1989.

This is the southernmost locality for the species. *Myrmecina* CURTIS, 1829 has its center of distribution in the Oriental region (EMERY, 1921; MANN, 1919, 1921; BROWN, 1973; ONOYAMA, 1980) with a single Australian species (TAYLOR & BROWN, 1985). The small *graminicola* group reaches Eurasia, North-America (BROWN, 1967; BERNARD, 1968) and Mediterranean Africa (CAGNIANT, 1973). The genus seems to be absent from the Ethiopian region (BOLTON, 1973). Some 35 taxa have been described; the comments of BERNARD (1968) about two hundred species seem without fundament. The genus has never been subjected to a formal revision.

* Departamento de Biología Animal, Biología Vegetal e de Ecología. Universidade Autónoma de Barcelona. 08193 Bellaterra (Spain)

** Departamento de Zoología. Universidade de La Laguna. 38206 La Laguna-Tenerife (Spain)

Myrmecina graminicola (LATR., 1802) and *M. sicula* (ANDRÉ, 1882) are the two single european species but this last taxon, a sicilian endemism, is probably a synonym (MENOZZI, 1936; BARONI URBANI, 1964). Typical biotopes for *M. graminicola* are humid places with a developed layer of humus. The nest is hypogaeous but occasionally workers can be seen foraging among dead leaves (pers. obs.). Feeding habits seem to be variable: in northern Europe it is a scavenging ant (COLLINGWOOD, 1979) and does not maintain nest aphids; according to BERNARD (1968) it is nectarivorous and can have nest aphids; dead or live animal food, specially insects, is the preferred regime as stated by DONISTHORPE (1927) but accepts also honey. FOREL (1920) also observed some workers taking honey in a lab nest and mentions a single worker licking a decaying cat carcass. Swarming takes place from August to October in North Europe (COLLINGWOOD, 1979), from August to September in Switzerland (KUTTER, 1977) and Spain (pers. obs., X.E.). No data are known for North Africa.

ACKNOWLEDGEMENTS

This work was supported by a grant ("Estudios sobre invertebrados terrestres de Madeira e Canarias") from the University of La Laguna, Tenerife to M. Báez.

REFERENCES

BARONI URBANI, C. :

1964. Studi sulla mirmecofauna d'Italia. II. Formiche di Sicilia. *Atti Academia Gioenia Scienze Naturali di Catania*, Ser. 6 (16):25-66.

BERNARD, F. :

1968. *Les fourmis (Hymenoptera, Formicidae) d'Europe occidentale et septentrionale*. Masson. Paris. pp. 411.

BOLTON, B. :

1973. The ant genera of West Africa: a synonymic synopsis with keys (Hymenoptera: Formicidae). *Bulletin of the British Museum of natural History (Entomology)*, 27: 319-368.

BROWN, W.L. :

1973. Studies on North American ants. II. *Myrmecina*. *Entomological News*, 78: 233-240.

BROWN, W.L. :

1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. In: *Tropical forest ecosystems in Africa and South America: A comparative review*: 161-185. (Meggers, B.J., Ayensu, E.S. & Duckworth, W.D. eds.). Smithsonian Institution Press. Washington.

CAGNIANT, H. :

1973. Les peuplements de fourmis des forêts algeriennes. *Ecologie. Biocénétique*, essai biologique. Ph. D. Thesis. Université Paul Sabatier, Toulouse. pp.464.

COLLINGWOOD, C.A. :

1979. The Formicidae (Hymenoptera) of Fennoscandia and Denmark. *Fauna Ent. Scandinavica*, Vol. 6. pp. 174.

- DONISTHORPE, H.J.K. :
1927. British ant. George Routeledge & Sons. London. pp. 436.
- EMERY, C. :
1921. Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae. In: *Genera Insectorum*. Fasc. 174 (Wytzman, p. ed.). Bruxelles.
- FOREL, A. :
1920. Les fourmis de la Suisse. Le Flambeau. La Cahux-de-Fonds. pp. 333.
- HEINZE, J. :
1986. A new find of *Paratrechina longicornis* (Latreille, 1802) on the Azores (Hymenoptera, Formicidae). *Bocagiana*, **101**: 1-3.
- KUTTER, H. :
1977. Hymenoptera. Formicidae. *Insecta Helvetica*, **6**. Zurich. pp. 289.
- MANN, M.M. :
1919. The ants of the British Solomon Island. *Bulletin of the Museum of Comparative Zoology*. **63**: 273-391.
- MENOZZI, C. :
1936. Nuovi contributi alla conoscenza della fauna delle isole italiane dell'Egeo. VI. Hymenoptera-Formicidae. *Bolletino del Laboratorio di Zoologia Generale e Agraria di Portici*, **29**: 262-311.
- ONOHAMA, K. :
1980. An introduction to the ant fauna of Japan, with a check list (Hymenoptera, Formicidae). *Kontyû*, **48**: 193-212.
- TAYLOR, R.W. & BROWM, D.R. :
1985. Hymenoptera: Formicoidea. *Zoological Catalogue of Australia*, Vol. 2: 1-149. Australia Gov. Pub. Serv. Canberra.
- TAYLOR, R.W. & WILSON, E.O. :
1961. Ants from three remote oceanic islands. *Psyche*, **68**: 137-144.
- WELLENIUS, O.H. :
1949. Die Formiciden von den Azoren und Madeira. *Societas Scientiarum Fennica. Commentationes Biologicae*, **8** (19) : 1-4.
- WHEELER, W.M. :
1927. The ants of the Canary Islands. *Proceedings of the American Academy of Arts and Sciences*, **62** : 93-210.
- WILSON, E.O. :
1973. The ants of Easter Islands and Juan Fernández. *Pacific Insects*, **15** : 285-287.