

According to Brown (1949a: 17) "this species ... will probably be found on any really tropical shore in the central and western part of the Pacific". The specimen from the Seychelles is not available from the main museum collections (Bolton, pers. com.).

Strumigenys rogeri Emery

Strumigenys rogeri Emery, 1890: 68.

Distribution. - Africa, America, Asia, Europe; SEYCHELLES: Cousin [G. M. Bathe].

Ecology. - Tramp species, probably of West African origin; nests normally in rotten wood (Bolton, 1983: 387ff).

Strumigenys scotti Forel

Strumigenys scotti Forel, 1912: 159.

Distribution. - Africa; SEYCHELLES: S: Silhouette (Mare aux Cochons, 305 m [type locality, H. M. Scott]).

Ecology. - Probably afrotropical species of limited tramping ability (Bolton, 1983).

Terataner scotti (Forel)

Atopomyrmex scotti Forel, 1912: 160.
Terataner scotti (Forel); Emery, 1912: 103.

Distribution. - Endemic species of the SEYCHELLES: S: Praslin [type locality, H. M. Scott], Silhouette (plateau de la Mare aux Cochons, 305 m [H. M. Scott], forest, 610 m [H. M. Scott]).

Ecology. - Unknown. *Terataner* species are strictly arboreal, nesting in the trunks or branches of living trees (Bolton, 1973).

Tetramorium bicarinatum (Nylander)

Myrmica bicarinata Nylander, 1846b: 1061.
Tetramorium guineense Fabricius, 1793: 357 *sensu* Mayr, 1862: 740 and all subsequent authors [misidentification, see Bolton, 1977: 94].

Tetramorium bicarinatum (Nylander); Bolton, 1977: 94.
[revisions of the genus *Tetramorium* in Bolton (1976, 1977, 1979, 1985)].

Distribution. - America, Asia, Australia, Europe, Malagasy Region; SEYCHELLES: S: Fregate [M. Mühlenberg; U. Müller], Grande Soeur [M. Mühlenberg], Petite Soeur [M. Mühlenberg].

Ecology. - Tramp species.

Tetramorium lanuginosum Mayr

Tetramorium lanuginosum Mayr, 1870: 972+976.
Tetramorium obesum *st. striatidens* Emery, 1889: 501 [syn. Bolton, 1976: 350].
Triglyphothrix striatidens var. *felix* Forel, 1912: 160 [syn. Bolton, 1976: 350].
Triglyphothrix striatidens (Emery); Bolton, 1976: 350 [syn. with *Triglyphothrix lanuginosa*].
Triglyphothrix lanuginosa Mayr; Bolton, 1985 [generic synonymy].

Distribution. - America, Asia, Australia, Malagasy Region; SEYCHELLES: AL: Aldabra (South Island [B. Cogan & A. Hutson]); S: Aride [J. Rowley & S. Warman], Félicité [H. M. Scott], Mahé (Union Vale [I. H. & J. B. Haines]), Silhouette (Mare aux Cochons [H. M. Scott]).

Ecology. - Tramp species; one single specimen collected on Aride in 1975, but none in 1978 (Rowley & Warman, 1979).

Tetramorium quadrispinosum Emery

Tetramorium quadrispinosum Emery, 1886: 362.
Tetramorium blochmannii var. *montanum* Forel, 1891: 153 [syn. Bolton, 1979: 155].
Tetramorium blochmannii (sic) *montanum* Forel; Emery, 1922: 282.
Tetramorium quadrispinosum *st. montanum* Forel; Santschi, 1918: 122; Emery, 1922: 282.

Distribution. - Africa, Malagasy Region; SEYCHELLES: AL: Aldabra [J. C. F. Fryer].

Ecology. - fairly common on Aldabra (Bolton, 1979: 155).