

*Ecology.* - Unknown.

***Camponotus (Tanaemyrmex) maculatus-gouldi-group***

*Camponotus (Tanaemyrmex) maculatus-gouldi-group* Emery, 1925: 84.

*Camponotus (Tanaemyrmex) maculatus-group* Emery *sensu* Haines & Haines, 1978a: 635.

*Distribution.* - Mahé ([I. H. & J. B. Haines], Union Vale [I. H. & J. B. Haines]).

*Ecology.* - See species above.

***Paratrechina bourbonica bourbonica* (Forel)**

*Prenolepis nodifera r. bourbonica* Forel, 1886a: 210.  
*Prenolepis bourbonica r. bengalensis* Forel, 1894a: 406+408 [syn. Wilson & Taylor, 1967a: 88].

*Prenolepis bengalensis* Forel; Bingham, 1903: 326.  
*Paratrechina (Nylanderia) bourbonica bengalensis* (Forel); Emery, 1925: 219.

*Distribution.* - Asia; Malagasy Region; SEYCHELLES: C: Coetivy [J. S. Gardiner]; F: St. Pierre [M. Mühlenberg]; S: Albatross [M. Mühlenberg], Aride [M. Mühlenberg], Chauve Souris [M. Mühlenberg], Cousin [M. Mühlenberg], Cousine [M. Mühlenberg], Fregate [M. Mühlenberg], Grande Soeur [M. Mühlenberg], La Digue [C. Alluaud], Mahé [C. Alluaud] (Cascade Estate [H. M. Scott], Petite Soeur [M. Mühlenberg], Praslin [M. Mühlenberg], ?Round [C. Alluaud]).

*Ecology.* - Tramp species; males and females were collected in December and January (Forel, 1912). Brown (1959) found the species in the Solomon Islands in coconut plantations.

***Paratrechina bourbonica farquharensis* (Forel)**

*Prenolepis bourbonica var. farquharensis* Forel, 1907a: 92.

*Paratrechina (Nylanderia) bourbonica var. farquharensis* (Forel); Emery, 1925: 219.

*Distribution.* - Subspecies: endemic of the SEYCHELLES: F: Farquhar [type locality, J. S. Gardiner].

*Ecology.* - Unknown.

***Paratrechina madagascariensis sechellensis* (Emery)**

*Prenolepis madagascariensis var. sechellensis* Emery, 1894a: 71.

*Paratrechina (Nylanderia) ellisi madagascariensis var. sechellensis* (sic) (Emery); Emery, 1925: 219 [unavailable name].

*Distribution.* - Species: endemic of the Malagasy Region; subspecies: endemic of the SEYCHELLES: S: Mahé [type locality of subspecies, C. Alluaud].

*Ecology.* - Unknown.

***Paratrechina mixta* (Forel)**

*Prenolepis mixta* Forel, 1897: 203.

*Paratrechina (Nylanderia) mixta* (Forel); Emery, 1925: 219.

*Distribution.* - Endemic species of the SEYCHELLES: S: ? [type locality, A. Brauer], Mahé [H. M. Scott], Praslin [H. M. Scott], Silhouette (forest above 610 m [H. M. Scott]).

*Ecology.* - Males and females were collected in August and September (Forel, 1912).

***Paratrechina steinheili* (Forel)**

*Prenolepis steinheili* Forel, 1893: 342.

*Paratrechina (Nylanderia) steinheili* (Forel); Emery, 1925: 223.

*Distribution.* - America; SEYCHELLES: AL: Aldabra [J. C. F. Fryer].

*Ecology.* - Species probably introduced from Central America (Emery, 1925: 223); biological data only from outside the Seychelles: Forel (1893) reported *P. steinheili* from the Antilles, where it is common, mainly in open places below 152 m, but also found in second growth and in shady places in forests up to 366 m; nest below a stone (even below coral blocks in thickets near the sea) or a log, under the bark of rotten logs, in a rotten stump or at