

- Legs almost uncolored.....	12
12 (11). Frontal lobe extremely developed, extending beyond anterior margin of clypeus (Fig. 53A, B).....	<i>P. upeneci</i> Forel [p.125]
- Frontal lobe sometimes developed, but never extending beyond anterior margin of clypeus (Figs. 50B, 52B).....	13
13 (12). Eye consisting of 5-7 ommatidia (Fig. 33B); terminal segment of antenna more than 1.6 times as long as preceding two segments together.....	<i>P. parvicorpus</i> sp. nov. [p.86]
- Eye consisting of at least 10 ommatidia; terminal segment of antenna at most 1.3 times as long as preceding two segments together.....	14
14 (13). Frons irregularly reticulate; hypostoma bearing a stout median process (Fig. 7C); frontal carina horizontal, extensively overhanging antennal scrobe (Fig. 7A); each dorsolateral portion of promesonotal dome produced outward (Fig. 7D).....	<i>P. aristotelis</i> Forel [p.34]
- Frons longitudinally rugose; hypostoma bearing 0-3 median processes; the condition of frontal carina and antennal scrobe variable; each dorsolateral portion of promesonotal dome sometimes produced outward.....	15
15 (14). Masticatory margin of mandible having two denticles before basal angle; ventral faces of midcoxa and hindcoxa distinctly reticulate (this condition exclusively shared with <i>P. aristotelis</i> among others).....	16
- Masticatory margin of mandible having at most one denticle before basal angle; ventral faces of midcoxa and hindcoxa completely smooth and shining.....	20
16 (15). Lateral portion of clypeus developed into a horn (Fig. 11).....	<i>P. clypeocornis</i> sp. nov. [p.44]
- Lateral portion of clypeus never developed into a horn (Figs. 20B, 23B, 43B, 51B).....	17
17 (16). Outer face of mandible sculptured with rugulae running from its base toward its masticatory margin (Fig. 43A)	<i>P. rugifera</i> sp. nov. [p.106]
- Outer face of mandible smooth and shining, or rugose laterally only around its base.....	18
18 (17). Frons (the area between frontal carinae) reticulate in its posterior part where spaces enclosed by rugulae punctured and dull.....	<i>P. tenebricosa</i> sp. nov. [p.121]
- Frons longitudinally rugose or rugoso-reticulate in its posterior part where spaces enclosed by rugulae almost smooth and shining.....	19
19 (18). Frons in profile relatively flat (Fig. 20B).....	<i>P. hortensis</i> Forel [p.61]
- Frons in profile relatively convex (Fig. 23B).....	<i>P. kikutai</i> sp. nov. [p.67]
20 (15). Frontal carina horizontal, extensively overhanging antennal scrobe (Fig. 52A); hypostoma always bearing one large median process (Fig. 52C).....	21
- Frontal carina sometimes well developed, but not overhanging antennal scrobe extensively; hypostoma bearing 0-3 median processes.....	23
21 (20). Clypeus having a distinct median longitudinal carina (Fig. 42A).... <i>P. retivertex</i> sp. nov. [p.104]	
- Clypeus lacking a median longitudinal carina.....	22
22 (21). Frontal lobe well developed and suberect (Fig. 52B); frons sparsely covered with longitudinal rugulae.....	<i>P. tjibodana</i> Forel [p.123]
- Frontal lobe poorly developed (Fig. 47B); frons densely covered with longitudinal rugulae.....	<i>P. sayapensis</i> sp. nov. [p.114]