

- 33 (30). Posterior declivity of promesonotal dome having a distinct prominence (Fig. 36D).....
 *P. plinii* Forel [p.92]
 - Posterior declivity of promesonotal dome lacking a distinct prominence..... 34
- 34 (33). Hypostoma bearing an indistinct median process, or lacking median process; clypeus having a median longitudinal carina (Fig. 26E)..... *P. lucioccipitalis* sp. nov. [p.73]
 - Hypostoma bearing 2-3 median processes; clypeus lacking a conspicuous median longitudinal carina..... 35
- 35 (34). Promesonotum narrower dorsally than ventrally..... *P. deltea* sp. nov. [p.47]
 - Promesonotum broader dorsally than ventrally..... *P. merimbun* sp. nov. [p.79]
- 36 (29). Posterior declivity of promesonotal dome lacking a distinct prominence; if the declivity has an inconspicuous prominence (as seen in *P. poringensis* sp. nov.), hypostoma bearing 2-3 distinct median processes..... 37
 - Posterior declivity of promesonotal dome having a distinct prominence; if the declivity has just an inconspicuous prominence (as seen in *P. submonticola* sp. nov.), hypostoma completely lacking median processes..... 43
- 37 (36). Frontal lobe relatively large and erect (Fig. 50B)..... *P. tawauensis* sp. nov. [p.119]
 - Frontal lobe relatively small (Figs. 8B, 9B, 35B)..... 38
- 38 (37). Posterior margin of head weakly concave in full-face view (Fig. 35A); dorsum of promesonotum smooth and shining (with several rugulae)..... *P. planidorsum* sp. nov. [p.91]
 - Posterior margin of head strongly concave in full-face view; dorsum of promesonotum sculptured or smooth..... 39
- 39 (38). Hypostoma bearing three median processes (Figs. 8C, 9C)..... 40
 - Hypostoma bearing a pair of stout median processes; if hypostoma bears three median processes, then the medianmost one is much reduced (as seen in *P. rabo* (Fig. 41C) and *P. poringensis* sp. nov.)..... 41
- 40 (39). Petiolar node relatively high (Fig. 8E)..... *P. bugi* Emery [p.37]
 - Petiolar node relatively low (Fig. 9D)..... *P. butteli* Forel [p.39]
- 41 (39). Petiole almost twice as long as postpetiole (excluding helcium) (Fig. 18D); lateral face of promesonotum almost smooth and shining..... *P. gombakensis* sp. nov. [p.57]
 - Petiole at most 1.7 times as long as postpetiole (excluding helcium) (Fig. 41D); lateral face of promesonotum sculptured..... 42
- 42 (41). Petiole ca. 1.7 times as long as postpetiole (excluding helcium) (Fig. 41D); outer face of mandible sparsely covered with very short appressed hairs..... *P. rabo* Forel [p.102]
 - Petiole 1.3-1.4 times as long as postpetiole (excluding helcium) (Fig. 37C); outer face of mandible sparsely covered with relatively long decumbent hairs..... *P. poringensis* sp. nov. [p.94]
- 43 (36). Hypostoma bearing three low median processes, of which each lateral one is partly combined with the process just mesal to mandibular insertion (Fig. 6C)..... *P. annexus* sp. nov. [p.32]
 - If hypostoma bears 2-3 median processes, then each lateral one is never combined with the process just mesal to mandibular insertion..... 44
- 44 (43). Hypostoma bearing a pair of stout median processes (Fig. 17C); petiole almost as long as postpetiole which is relatively massive (Fig. 17D)..... *P. ghigii* Emery [p.55]
 - If hypostoma bears a pair of small or indistinct median processes, then petiole is at least 1.2 times