

- than 1.2 times as long as postpetiole..... 31
- Hypostoma of the major bearing 1-3 median processes (in some specimens of *P. lucioccipitalis* sp. nov. median processes sometimes almost absent where petiole is more than 1.5 times as long as postpetiole in both the subcastes)..... 33
- 31 (30). Distance between mandibular insertion and anterior margin of eye 2.3-2.4 times as long as maximal diameter of eye in the major (Fig. 19B), and ca. 1.5 times in the minor.....  
..... *P. havilandi* Forel [p.59]
- Distance between mandibular insertion and anterior margin of eye 1.4-1.5 times as long as maximal diameter of eye in the major, and 1.0-1.1 times in the minor..... 32
- 32 (31). Postpetiole 2.4-2.6 times as broad as petiolar node in the major; frontal carina of the major conspicuous..... *P. manukana* sp. nov. [p.75]
- Postpetiole 1.8-1.9 times as broad as petiolar node in the major; frontal carina of the major inconspicuous..... *P. megacephala* (Fabricius) [p.77]
- 33 (30). Posterior declivity of promesonotal dome having a distinct prominence in both the subcastes (Fig. 36D, F); promesonotal dome of the minor having a pair of low tubercles (Fig. 36F).....  
..... *P. plinii* Forel [p.92]
- Posterior declivity of promesonotal dome lacking a distinct prominence in both the subcastes; promesonotal dome of the minor unarmed..... 34
- 34 (33). Hypostoma of the major bearing an indistinct median process, or lacking median processes; clypeus of the major having a median longitudinal carina (Fig. 26A)... *P. lucioccipitalis* sp. nov. [p.73]
- Hypostoma of the major bearing 2-3 median processes; clypeus of the major lacking a median longitudinal carina..... 35
- 35 (34). Promesonotum of the major narrower dorsally than ventrally; propodeal spine of the minor 3.5-4 times as long as diameter of propodeal spiracle (Fig. 13E)..... *P. deltea* sp. nov. [p.47]
- Promesonotum of the major broader dorsally than ventrally; propodeal spine of the minor twice as long as diameter of propodeal spiracle (Fig. 29E)..... *P. merimbun* sp. nov. [p.79]
- 36 (29). Posterior declivity of promesonotal dome of the major lacking a distinct prominence; if the declivity has an inconspicuous prominence (e.g., *P. poringensis* sp. nov.), hypostoma of the major bears 2-3 distinct median processes..... 37
- Posterior declivity of promesonotal dome of the major having a distinct prominence; if the declivity has just a small prominence as seen in *P. submonticola* sp. nov. (Fig. 49C), hypostoma of the major completely lacks median processes..... 43
- 37 (36). Promesonotum of the minor smooth and shining..... 38
- Promesonotum of the minor sculptured..... 41
- 38 (37). Promesonotal dome of the minor in profile very low and almost flat dorsally (Fig. 35F); in full-face view posterior margin of head of the major weakly concave (Fig. 35A).....  
..... *P. planidorsum* sp. nov. [p.91]
- Promesonotal dome of the minor in profile relatively convex dorsally; in full-face view posterior margin of head of the major usually distinctly concave..... 39
- 39 (38). Hypostoma of the major with a pair of very large median processes (Fig. 18C); dorsum of head of the minor rugoso-reticulate..... *P. gombakensis* sp. nov. [p.57]
- Hypostoma of the major with 2-3 median processes; if two processes are present, they are much