

- smaller than those of *P. gombakensis* sp. nov. (Fig. 18C); head of the minor largely smooth and shining..... 40
- 40 (39). Frontal lobe of the major relatively large and erect (Fig. 50B)... *P. tawauensis* sp. nov. [p.119]
- Frontal lobe of the major relatively small (Fig. 9B)..... *P. butteli* Forel [p.39]
- 41 (37). Outer face of mandible of the major bearing long decumbent hairs (Fig. 37B); head of the minor reticulate..... *P. poringensis* sp. nov. [p.94]
- Outer face of mandible of the major sparsely bearing short appressed hairs; head of the minor punctured..... 42
- 42 (41). Distance between mandibular insertion and anterior margin of eye twice as long as maximal diameter of eye in the major (Fig. 41B); hypostoma of the major bearing three median processes, of which medianmost one is poorly developed (Fig. 41C)..... *P. rabo* Forel [p.102]
- Distance between mandibular insertion and anterior margin of eye ca. 1.4 times as long as maximal diameter of eye in the major (Fig. 8B); hypostoma of the major bearing three similar-sized median processes (Fig. 8C)..... *P. bugi* Wheeler [p.37]
- 43 (36). Hypostoma of the major bearing three low median processes, of which each lateral one is partly connected to the process just mesal to mandibular insertion (Fig. 6C)..... *P. annexus* sp. nov. [p.32]
- If hypostoma of the major bears 2-3 median processes, each lateral one is never connected to the process just mesal to mandibular insertion..... 44
- 44 (43). Hypostoma of the major bearing a pair of stout median processes (Fig. 17C); petiole of the major almost as long as postpetiole (Fig. 17D); promesonotum with a pair of low tubercles in the minor..... *P. ghigii* Emery [p.55]
- If hypostoma of the major bears a pair of small or indistinct median processes, petiole of the major is at least 1.2 times as long as postpetiole; promesonotum without a pair of low tubercles in the minor..... 45
- 45 (44). Propodeal spine of the major elongate-triangular, with broad base (Fig. 10C)..... 46
- Propodeal spine of the major horn-like, with narrow base (Fig. 4D)..... 47
- 46 (45). Antennal scrobe margined below with a conspicuous carina in the major anteriorly (Fig. 10A); mesopleuron of the major without a distinct transverse impression; posterior declivity of promesonotal dome of the minor with a low prominence (Fig. 10F).... *P. cariniceps* sp. nov. [p.41]
- Antennal scrobe present only around antennal insertion, and never margined below with a conspicuous carina in the major (Fig. 49A); mesopleuron of the major divided by a transverse impression into two parts, of which the lower part is margined dorsally; posterior declivity of promesonotal dome of the minor without any prominence (Fig. 49E).....
- *P. submonticola* sp. nov. [p.117]
- 47 (45). Posteriormost part of dorsum of occipital lobe of the major almost transversely rugose in the major (Fig. 34A)..... *P. plagiaria* Fr. Smith [p.88]
- Posteriormost part of dorsum of occipital lobe reticulate, rugoso-reticulate or longitudinally rugose, but never transversely rugose in the major..... 48
- 48 (47). Frontal carina of the major slight; lateral face of occipital lobe of the major smooth or very weakly rugoso-punctured..... *P. orophila* sp. nov. [p.85]
- Frontal carina of the major distinct; lateral face of occipital lobe of the major rugoso-reticulate..... 49
- 49 (48). Posterior margin of head of the major strongly concave (Fig. 4A, B); occipital carina of the minor