

0.48, EL 0.12-0.13, WL 0.64-0.66, PW 0.16-0.17, PL 0.13-0.14, PPW 0.24-0.25, PPL 0.15-0.16. Indices: CI 76-78, SI 76-77, PI 114-131, PPI 156-160.

Mandibles with 5 teeth; anterior border of clypeus convex, clypeal surface convex, with carinae (including middle carina) poorly defined; vertex concave; eyes small, with 7 or 8 ommatidia in greatest diameter, maximum diameter usually less than shortest distance to insertion of mandible; scape barely reaching occipital corner; mesosoma without sutures depressed below surrounding area; propodeal spines consisting of tiny angles; petiole with blunt node as seen in profile (Fig. 100), anterior petiolar face slightly concave, posterior face convex, subpeduncular process well formed, with blunt, rounded tip.

Erect hairs sparse, but present on most surfaces, short (most < 0.05); decumbent pubescence obvious only on head, and consisting of few scattered hairs.

Sculpture of head fine, dorsum nearly completely smooth and shining, but with a few fine striae on sides, striae well defined anterior to eyes and near frontal carinae, mesosoma, petiole and postpetiole punctate, side of pronotum with striae, gaster smooth and shining.

Color: light yellow brown.

Female measurements (mm): HL 0.74, HW 0.64, SL 0.56, EL 0.22, WL 1.13, PW 0.23, PL 0.16, PPW 0.38, PPL 0.19. Indices: CI 86, SI 76, PI 144, PPI 200.

Similar to worker in most aspects, differing in that the eye is larger, diameter about 0.7 times minimum distance to insertion of mandibles; ocelli well developed; scape reaching occipital corner; vertex concave; mesosoma large; propodeal spines well developed, length about 0.8mm; subpeduncular process well developed, rounded tip; petiole similar to that of worker, except node narrower.

Hairs as in worker, blunt tipped, some weakly spatulate, those on gaster up to 0.10mm in length; decumbent pubescence nearly absent, few hairs on head and gaster.

Sculpture slightly more coarse than in worker, dorsum of head completely striate, surfaces lateral to and anterior to eye nearly rugose, scutum and scutellum finely striate, side of pronotum and mesopleuron finely striate, propodeum rugose and punctate in intrarugal spaces, petiole and postpetiole punctate, with rugae on anterior face of petiole, transversely across top of node, on side and across posterior half of postpetiole, gaster smooth and shining.

Color: yellow brown, slightly darker than workers.

Male: Unknown.

Type series: Holotype worker (MCZC), 21 paratype workers (AMNH,