(Atkinson et al. 1991). There are no native cockroaches in western Canada today (Kevan 1979). These and other amber insects indicate a subtropical to tropical climate in the Pacific Northwest some 50–55 million years ago, which is supportive of previously derived conclusions from analyses of fossil plants assemblages from this period and locality (Verschoor 1974; Hopkins 1980; Wilson 1996).

Acknowledgments

We would like to thank Mark Winston and the Biology Department, Simon Fraser University, for financial assistance, The British Columbia Paleontological Alliance for providing a Research Site Reserve permit, Mayumi Kariya of British Columbia Hydro for granting access permission, Dave Langevin for providing specimens for this study, J.B. Lambert and C.E. Shawl for the NMR spectra, and Shelley Dowson, Mark Winkler, and Brain Jackson for assistance at the site. We thank C. Baroni Urbani for determining the Hat Creek ant specimens and supplying information from his cladistic analysis of the Apocrita concluding that the New Jersey "ant" fossils are much closer to wasps than to true ants of the family Formicidae.

References

- Agosti D, Grimaldi D, Carpenter JM. 1998. Oldest known ant fossils discovered. Nature (London) 391: 447. Atkinson TH, Koehler PG, Patterson RS. 1991. Catalog and atlas of the cockroaches (Dictyoptera) of North America north of Mexico. Miscellaneous Publications of the Entomological Association of America 78.
- Baroni Urbani C. 1989. Phylogeny and behavioral evolution in ants. Ethology, Ecology and Evolution 1: 137-68.
- Bell WA. 1957. Flora of the Upper Cretaceous Nanaimo Group, Vancouver Island. Geological Survey of Canada Memoir 293.
- Brandao CRF, Baroni Urbani C, Wagensberg J, Yamamoto CI. 1999. New *Technomyrmex* in Dominican amber (Hymenoptera: Formicidae), with a reappraisal of the Dolichoderinae phylogeny. *Entomologica Scandinavica*. In press.
- Carpenter FM. 1929. A fossil ant from the Lower Eocene (Wilcox) of Tennessee. Journal of the Washington Academy of Science 19: 300-301.
- 1930. The fossil ants of North America. Bulletin of the Museum of Comparative Zoology 70: 1-66.
- Cockerell TDA. 1921. Some Eccene insects from Colorado and Wyoming. Proceedings of the United States National Museum 59: 29-39.
- Church BN. 1975. Geology of the Hat Field Creek Coal Basin. pp. 99-118 in Fieldwork 1975. British Columbia Ministry of Energy, Mines and Petroleum Resources Paper 921/13E.
- 1981. Further studies of the Hat Creek coal deposit. pp. 73-77 in Geological Fieldwork 1980. British Columbia Ministry of Energy, Mines and Petroleum Resources Paper 1981-1.
- Church BN, Matheson A, Hora ZD. 1979. Combustion metamorphism in the Hat Creek area, British Columbia. Canadian Journal of Earth Sciences 16: 1882-87.
- Dlussky GM. 1975. Formicidae. pp. 115-121 in AP Rasnitsyn (Ed.), Hymenoptera Apocrita of the Mesozoic. Transactions of the Paleontological Institute of the Academy of Sciences of the USSR 147. [In Russian.]
- Grand L. 1984. Paleontology of the Green River Formation, with a review of the fish fauna. 2nd ed. Bulletin of the Geological Survey of Wyoming 63: 1-333.
- Grimaldi D, Agosti D, Carpenter JM. 1997. New and rediscovered primitive ants (Hymenoptera: Formicidae) in Cretaceous amber from New Jersey, and their phylogenetic relationships. American Museum Novitates 3208.
- Hong Y-C, Yang T-C, Wang S-T, Wang S-E, Li Y-K, Sun M-R, Sun H-C, Tu N-C. 1974. Stratigraphy and paleontology of Fushin Coal field, Liaoning Province. Acta Geologica Sinica 2: 113-49.
- Hopkins WS Jr. 1980. Palynology of the 75-106 B.C. Hydro core hole, Hat Creek Coal Basin, British Columbia. Geological Survey of Canada Open File 547.
- Johnston JE. 1993. Insects, spiders, and plants from the Tallahatta Formation (Middle Eocene) in Benton County, Mississippi. Mississippi Geology 14: 71-82.