



Fig. 3. *Cataglyphis* Foerster. (a, b) *C. foreli*, lectotype; (c, d) *C. oxianus*, paratype; (e) *C. ruber*; (f) *C. cuneinodis*; (g, h) *C. cinnamomeus*, syntype; (i-k) *C. albicans*; (l) *C. alibabae*, paratype. (a) Lateral view of thorax and petiole; (b, d) frontal view of head; (c) lateral view of propodeum and petiole; (e, f, k, l) lateral view of petiole; (g, i) 3rd-6th maxillary palpomeres; (h, j) 1st-3rd flagellomeres. All figures related to workers.

55(54). Petiole rather high, cuneiform; in lateral view, node higher than long (Fig. 3f). Occipital margin of head without erect hairs.—S Transcaucasia, Iran *C. cuneinodis* Arnoldi, stat. n.

56(53). Body uniformly brown to black.

57(60). Maxillary palps short; 6th palpomere nearly as long as 5th (Fig. 3g). 2nd flagellomere distinctly shorter than 3rd and 2-2.3 times as long as wide (Fig. 3h).

58(59). Head and large part of thorax completely smooth, strongly shining. Node of petiole longer than high (in lateral view).—Kopet Dag *C. elegantissimus* Arnoldi.

59(58). Body with fine but distinct superficial sculpture, weakly shining. Node of petiole as high as, or higher than long (in lateral view).—Middle

Asia, S Kazakhstan, Afghanistan *C. cinnamomeus* (Karawajew).

60(57). Maxillary palps long; 6th palpomere 0.6-0.7 times as long as 5th (Fig. 3i). 2nd flagellomere as long as 3rd and 3 times as long as wide (Fig. 3j).

61(62). Head and thorax more coarsely sculptured, semi-dull.—N Africa, Lebanon, Arabia *C. semitonsus* Santschi.

62(61). Head and thorax with smoothed sculpture; integument at least weakly shining.

63(64). Petiole low, broadly rounded or slightly flattened at apex (Fig. 3k).—Spain, N Africa, Sudan, Ethiopia, Middle East, Arabia, Transcaucasia, Afghanistan *C. albicans* (Röger).