

Mt. Deogyu-san, Is. Eocheong-do, Kunsan, Mt. Mai-san, Is. Seonyu-do, Is. Wi-do, Kimjae. [JN] Is. Jaeun-do, Mt. Weolchul-san, Mt. Yudal-san, Is. Bikeum-do, Is. Odong-do, Is. Hong-do, Is. Daeheugsan-do, Is. Soheugsando, Is. Keokeum-do, Is. Jindo, Is. Wan-do, Is. Soan-do, Is. Bokil-do. [GB] Is. Uleung-do. [GN] Is. Juk-do, Mt. Baegam-san, Masan, Jinhae, Keojae, Namhae, Is. Hansan-do, Is. Wonsan-do. [JJ] Mt. Hanla-san.

Distribution. Korea, Japan.

Remarks. Over all, this species is similar to the species of *Pyramica* in body size and shape, but differ in having thin, elongate mandibles rather than short and triangular ones.

Genus *Pyramica* Roger, 1862

Pyramica Roger, 1862, Berl. Entomol. Z. 6: 251. Type species: *Pyramica gundlachi* Roger, 1862: 253, by monotypy.

Cephaloxys Smith, F. 1865, J. Proc. Linn. Soc. Lond. Zool. 8: 76. Type species: *Cephaloxys capitata* Smith, F. 1865: 77, by monotypy.

Glamyromyrmex Wheeler 1915, Bull. Mus. Comp. Zool. 59: 487. Type species: *Glamyromyrmex beebei* Wheeler, 1915: 488, by monotypy.

Smithistruma Brown 1948, Trans. Am. Entomol. Soc. 74: 104. Type species: *Strumigenys pulchella* Emery, 1895: 327.

Epitritus Emery, 1869, Bull. Soc. Entomol. Ital. 1: 136. Type species: *Epitritus argiolus* Emery, 1869: 136, by monotypy.

Kyidris Brown, 1949b, Mushi 20: 3. Type species: *Kyidris mutica* Brown, 1949b: 3.

Miccostruma Brown, 1948, Trans. Am. Entomol. Soc. 74: 123. Type-species: *Epitritus mandibularis* Szab, 1909: 27.

Platystruma Brown, 1953, Am. Midl. Nat. 50: 112. Type-species: *Strumigenys (Cephaloxys) depressiceps* Weber, 1934: 47.

Polyhomoa Azuma, 1950, Hyogo Biology 4: 36. Type-species: *Polyhomoa itoi* Azuma, 1950: 36, by monotypy.

***Pyramica incerta* (Brown, 1949) 긴털 톱비늘개미**

(Figs 9–10)

Smithistruma incerta Brown, 1949b, Mushi 20: 10; Choi, 1995: 191; Choi and Lee, 1995: 193; Choi, 1996a: 209; Choi, 1996c: 47; Kim, 1996: 181; Choi, 1997a: 57.

Smithistruma habei Onoyama, 1980: 194.

Pyramica incerta: Bolton, 1999, J. Nat. Hist. 33: 1673.

Workers. Body length around 1.5 mm. Body color yellowish to reddish brown. Lateral margins of head roundly convex. Clypeus fringed with spatulate hairs, and with a slightly concave anterior margin. Antennae 6 segmented. Leading edges of antennal scapes angulate at the basal third. Promesonotum more or less convex. Dorsal surface of mesosoma with numerous standing hairs. Pronotal humeri with elongate paired flagellate hairs. Sides of petiole and postpetiole and lower part of the gaster with masses