

record from the western Queensland Gulf Country (Lyrian Downs, N of Julia Creek), well outside the main distribution and the inland-most record of the genus in Australia. It is sympatric with *D. australe* in the area south of Cooktown. While it is currently unknown from Papua New Guinea (based on material examined during this study) it may well occur there as these two countries share many species across a range of taxonomic groups.

### Acknowledgements

We wish to dedicate this paper to Stefan Schödl. Stefan spent several weeks working in the Australian National Insect Collection as part of his studies on Australian *Meranoplus*. Unfortunately anticipated return visits were not realised. For invaluable assistance during this study we would like to thank Bernhard Merz (Département d'Entomologie, Muséum d'histoire naturelle, Geneva) for notes on and images of Forel material held in his care, and George Else and Kim Goodger (The Natural History Museum, London) for notes on and images of Fabricius specimens deposited in London. Valuable comments on the manuscript were received from Alan Andersen, Brian Heterick, Herbert Zettel, an anonymous reviewer, and the journal's editors. Funding for this study was generously provided by CSIRO Entomology.

### Zusammenfassung

Die australischen Arten der Ameisengattung *Diacamma* (Unterfamilie Ponerinae) werden revidiert. Vier Arten kommen in Australien vor: *D. australe* (FABRICIUS, 1775), *D. colosseense* FOREL, 1915, *D. leve* CRAWLEY, 1915 und *D. schoedli* sp.n. Ihre Vorkommen beschränken sich auf den Norden von Queensland und die Region Top End im Northern Territory. *Diacamma colosseense* und *D. leve* werden aus ihrer Synonymie mit *D. australe* gehoben. Für *D. colosseense* wird ein Lectotypus designiert.

### References

BILLEN, J.P.J. & PEETERS, C. 1991: Fine structure of the gemma gland in the ant *Diacamma australe* (Hymenoptera, Formicidae). – Belgian Journal of Zoology 121: 203-210.

BINGHAM, C.T. 1903: The Fauna of British India, including Ceylon and Burma. Hymenoptera. Ants and cuckoo-wasps. – Taylor & Francis, London, Vol. 2, 507 pp.

BITSCH, J. & PEETERS, C. 1991: Moignons alaires et morphologie thoracique chez l'ouvrière de la fourmi *Diacamma australe* (FABRICIUS) (Hym. Formicidae Ponerinae). – Bulletin de la Société Entomologique de France 96: 213-221.

BOLTON, B. 1995: A New General Catalogue of the Ants of the World. – Harvard University Press, Cambridge, Mass., 504 pp.

CRAWLEY, W.C. 1915: Ants from north and central Australia, collected by G. F. Hill. - Part I. – Annals and Magazine of Natural History (8)15: 130-136.

CUVILLIER-HOT, V., GADAGKAR, R., PEETERS, C. & COBB, M. 2002: Regulation of reproduction in a queenless ant: aggression, pheromones and reduction in conflict. – Proceedings of the Royal Society of London B 269: 1295-1300.

FABRICIUS, J.C. 1775: Systema entomologiae, sistens insectorum classes, ordines, genera, species adiectis synonymis, locis, descriptionibus, observationibus. – Korte, Flensburg and Leipzig, 832 pp.

FOREL, A. 1915: Results of Dr. E. Mjöberg's Swedish Scientific Expeditions to Australia 1910-13. 2. Ameisen. – Arkiv för Zoologi 9: 1-119.

HÖLLDOBLER, B. & ENGEL-SIEGEL, H. 1982: Tergal and sternal glands in male ants. – Psyche (Cambridge) 89: 113-132.

LE GUILLOU, E.J.F. 1842: Catalogue raisonné des insectes hyménoptères recueillis dans le voyage de circumnavigation des corvettes l'Astrolabe et La Zélée. – Annales de la Société Entomologique de France 10: 311-324.

MAYR, G.L. 1862: Myrmecologische Studien. – Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 12, Abhandlungen: 649-776.

PAUL, J. 2001: Mandible movements in ants. – Comparative Biochemistry and Physiology Part A 131: 7-20.

PEETERS, C. & BILLEN, J.P.J. 1991: A novel exocrine gland inside the thoracic appendages ('gemmae') of the queenless ant *Diacamma australe*. – Experientia 47: 229-231.

PEETERS, C., BILLEN, J.P.J. & HÖLLDOBLER, B. 1992: Alternative dominance mechanisms regulating monogyny in the queenless ant genus *Diacamma*. – Naturwissenschaften 79: 572-573.

PEETERS, C. & HIGASHI, S. 1989: Reproductive dominance controlled by mutilation in the queenless ant *Diacamma australe*. – Naturwissenschaften 76: 177-180.

RAMASWAMY, K., PEETERS, C., YUVANA, S.P., VARGHESE, T., PRADEEP, H.D., DIETEMANN, V., KARPAKAKUNJARAM, V., COBB, M. & GADAGKAR, R. 2004: Social mutilation in the Ponerine ant *Diacamma*: cues originate in the victims. – Insectes Sociaux 51: 410-413.

SHATTUCK, S.O. 1999: Australian ants. Their biology and identification. – Monographs on Invertebrate Taxonomy 3: 1-226.

SHATTUCK, S.O. & BARNETT, N.J. 2001: Australian Ants Online. – <<http://www.ento.csiro.au/science/ants/default.htm>>, retrieved on 1 April 2006.

TAYLOR, R.W. & BROWN, D.R. 1985: Formicoidea. Pp. 1-149, 306-348 in: WALTON, D.W. (ed.): Zoological catalogue of Australia, vol. 2. Hymenoptera: Formicoidea, Vespoidea and Sphecoidea. – Australian Government Publishing Service, Canberra, vi + 381 pp.

WILSON, E.O. 1958: Studies on the ant fauna of Melanesia. 3. *Rhytidoponera* in western Melanesia and the Moluccas. 4. The tribe Ponerini. – Bulletin of the Museum of Comparative Zoology at Harvard College 119: 303-371.

ZIMSEN, E. 1964: The type material of I. C. Fabricius. – Munksgaard, Copenhagen, 656 pp.