

## First record of *Pyramica baudueri* (EMERY, 1875) (Hymenoptera: Formicidae) from Bulgaria

Pavel BEZDĚČKA & Klára BEZDĚČKOVÁ



### Abstract

The first record of *Pyramica baudueri* (EMERY, 1875) from Bulgaria is reported. It also represents the discovery of a genus and tribe of ants unknown from Bulgaria until now. One worker of *P. baudueri* was found in South-eastern Bulgaria, near the village of Shiroko Pole. The known distribution of *P. baudueri* in the Palaearctic region is summarised and discussed.

**Key words:** Ants, first record, *Pyramica baudueri*, Dacetini, Formicidae, Bulgaria.

Myrmecol. News 13: 1-2

ISSN 1994-4136 (print), ISSN 1997-3500 (online)

Received 17 June 2009; revision received 21 July 2009; accepted 22 July 2009

Pavel Bezděčka (contact author) & Dr. Klára Bezděčková, Museum of Highlands Jihlava, Masarykovo nám. 55, CZ-586 01 Jihlava, Czech Republic. E-mail: bezdecka@muzeum.ji.cz

### Introduction

The Bulgarian myrmecofauna is insufficiently explored. In particular, some sub-Mediterranean species, parasitic species and species with cryptic lifestyle known from South-eastern Europe have not been reported from Bulgaria until now. In recent years several new records of ant species were reported for Bulgaria (see, e.g., RADCHENKO & ANTONOVA 2004, STANKIEWICZ & ANTONOVA 2005, STEINER & al. 2005, ANTONOVA 2008), but the last published checklist of Bulgarian ants (ATANASSOV & DLUSSKY 1992) has not been updated yet.

Here we report the discovery of the sub-Mediterranean species *Pyramica baudueri* (EMERY, 1875), a species with cryptic lifestyle, found by P. Bezděčka in South-eastern Bulgaria.

### Results

The recent species-level redetermination of ants deposited in the Museum of South-eastern Moravia (the Czech Republic, Zlín) resulted in the identification of one worker of *Pyramica baudueri* (EMERY, 1875). The redetermination of this specimen was done using the keys in HERVÉ (1969), AGOSTI & COLLINGWOOD (1987) and BOLTON (2000). The worker was collected in a place called Zhelezni vrata (41° 37' 06" N, 25° 28' 00" E, ca. 320 m a.s.l., South-eastern Bulgaria, Fig. 1) on 26.IV.1986 (leg. et det. P. Bezděčka). This forest-steppe locality lies two kilometres south of the centre of the village of Shiroko Pole. The specimen was found under a big stone on the soil surface. In spite of intensive exploration of the locality neither a nest nor further individuals were discovered.

### Discussion

*Pyramica* ROGER, 1862 is the second largest genus of the tribe Dacetini with 349 recent species (YUSUPOV 2009). Only five *Pyramica* species occur in the Western Palaearc-



Fig. 1: Habitat of *Pyramica baudueri* near Shiroko Pole (Photo: P. Bezděčka 1986).

tic region: *P. argiola* (EMERY, 1869), *P. baudueri*, *P. membranifera* (EMERY, 1869), *P. tenuipilis* (EMERY, 1915) and *P. tenuissima* (BROWN, 1953). The known distribution of *P. baudueri* covers the Channel Islands (RADCHENKO 2007), Morocco (BOLTON 2000), continental Spain (ESPADALER 1997, BOLTON 2000, RADCHENKO 2007), the Balearic Islands (ESPADALER 1997), Tunisia (BOLTON 2000), Malta (RADCHENKO 2007), continental France (BOLTON 2000, RADCHENKO 2007), Corsica (BOLTON 2000), Sardinia (BOLTON 2000, RADCHENKO 2007), Sicily (RADCHENKO 2007), continental Italy (BOLTON 2000, RADCHENKO 2007), Switzerland (BARONI URBANI 1998, BOLTON 2000, BRASCHLER 2002), Hungary (GALLÉ & al. 1998), Romania (MARKÓ 2008), Croatia (BRAČKO 2006), Macedonia (RADCHENKO 2007), Montenegro (PETROV 2006), Greece (AGOSTI & COLLINGWOOD 1987, RADCHENKO 2007), Turkey (BOLTON 2000, RADCHENKO 2007), and Armenia (ARAKELYAN 1994).

*Pyramica bauduerei* is exclusively a predator, hunting especially Collembola but occasionally also other small arthropods (MARKÓ 2008). It lives and forages in the soil, inhabiting sun-exposed and warm sites (BRASCHLER 2002, MARKÓ 2008). Colonies are usually small and monogynous (BOLTON 2000). Specimens of genus *Pyramica* can be identified as members of the tribe Dacetini immediately in the field, due to the characteristic shape of the head. A recent, detailed characterisation of workers of *P. bauduerei* is given by MARKÓ (2008).

The worker of *Pyramica bauduerei* reported here was found by hand collecting. The discovery represents the finding of a species, genus and tribe of ants hitherto unknown from Bulgaria. The hidden lifestyle of *Pyramica* ants and the hitherto lack of applying more specific collecting methods are probably the main reasons for the absence of any further data on this species from Bulgaria.

Recently, dacetine ants have been recorded more frequently in other parts of Europe, by using quantitative methods of collection. BRASCHLER (2002) caught two workers of *P. bauduerei* by pitfall trapping in Switzerland. DEVÁN (2008) captured the only specimen of the genus *Pyramica* known from Slovakia (male of *P. cf. argiola*) in a Moe-ricky trap (yellow dish with salt solution). FELLNER & al. (2009) repeatedly detected *P. argiola* in Austria using the Winkler sifting and extracting method, a collection technique commonly used in tropical rainforests. The latter two instances indicate that the application of a greater variety of collection techniques may boost European ant faunistics.

#### Acknowledgements

We thank the Ministry of Culture of the Czech Republic for the financial support of this research project (DE07PO40 MG009 MK ČR). We are very grateful to B. Braschler, P. Deván and A. Radchenko for valuable information and two anonymous referees for inspiring suggestions.

#### Zusammenfassung

Der erste Fund von *Pyramica bauduerei* (EMERY, 1875) in Bulgarien wird gemeldet. Es handelt sich dabei gleichzeitig um eine aus Bulgarien bisher unbekannte Gattung und Tribus von Ameisen. Eine Arbeiterin von *P. bauduerei* wurde im südöstlichen Bulgarien gefunden, nahe dem Dorf Shiroko Pole. Die bekannte Verbreitung von *P. bauduerei* in der Paläarktis wird zusammengefasst und diskutiert.

#### References

AGOSTI, D. & COLLINGWOOD, C.A. 1987: A provisional list of the Balkan ants (Hym., Formicidae) and a key to the worker caste. I. Synonymic list. – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 60: 51-62.

ANTONOVA, V. 2008: First record of the slave-maker ant *Harpagoxenus sublaevis* (NYLANDER, 1849) from Bulgaria (Hymenoptera: Formicidae). – Myrmecological News 12: 1-2.

ARAKELYAN, G.R. 1994: Fauna of the Republic of Armenia. Hymenopterous insects. Ants (Formicidae). – Gitutium, Erevan, 153 pp. (in Russian)

ATANASSOV, N. & DLUSSKY, G.M. 1992: Hymenoptera, Formicidae. – Fauna Bulgarica 22: 1-310.

BARONI URBANI, C. 1998: *Strumigenys bauduerei* (EMERY): espèce nouvelle pour la Suisse (Hymenoptera: Formicidae). – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 71: 163-164.

BOLTON, B. 2000: The ant tribe Dacetini. – Memoirs of the American Entomological Institute 65: 1-1028.

BRAČKO, G. 2006: Review of the ant fauna (Hymenoptera: Formicidae) of Croatia. – Acta Entomologica Slovenica 14: 131-156.

BRASCHLER, B. 2002: Neue Aspekte zur Verbreitung von *Pyramica bauduerei* (EMERY, 1875) (Hymenoptera, Formicidae). – Mitteilungen der Entomologischen Gesellschaft Basel 52: 139-142.

DEVÁN, P. 2008: The ants of Omšenská dolina valley. – Naturae Tutela 12: 101-104. (in Slovakian)

ESPADALER, X. 1997: Catàleg de les formigues (Hymenoptera: Formicidae) dels països catalans. – Sessió Conjuncta d'Entomologia ICHN-SCL 9: 23-42.

FELLNER, T., BOROVSKEY, V. & FIEDLER, K. 2009: First records of the dacetine ant species *Pyramica argiola* (EMERY, 1869) (Hymenoptera: Formicidae) from Austria. – Myrmecological News 12: 167-169.

GALLÉ, L., CSÓSZ, S., TARTALLY, A. & KOVÁCS, É. 1998: A checklist of Hungarian ants (Hymenoptera: Formicidae). – Folia Entomologica Hungarica 59: 213-220.

HERVÉ, P. 1969: Les espèces françaises de la tribu des Dacetini (Hymenoptera, Formicidae). – Entomops 13: 155-158.

MARKÓ, B. 2008: *Pyramica bauduerei* (EMERY, 1875) – a new ant species (Hymenoptera: Formicidae) for the Romanian fauna. – Fragmenta Faunistica 51: 101-106.

PETROV, I.Z. 2006: The ants of Serbia and Montenegro. – Serbian Academy of Sciences and Arts, Beograd, 136 pp. (in Serbian)

RADCHENKO, A. 2007: Fauna Europaea: Formicidae. In: NOYES, J. (Ed.): Fauna Europaea: Hymenoptera: Apocrita. – Fauna Europaea version 1.3; <<http://www.faunaeur.org>>, retrieved on 20 May 2009.

RADCHENKO, A. & ANTONOVA, V. 2004: First record of *Myrmica rugulososcabinodis* KARAWAJEW, 1929 (Hymenoptera: Formicidae) from Bulgaria. – Myrmecologische Nachrichten 6: 19-21.

STANKIEWICZ, A. & ANTONOVA, V. 2005: *Myrmica vandeli* BONDROIT (Hymenoptera: Formicidae) – a new ant species for Bulgaria. – Acta Zoologica Bulgarica 57: 123-126.

STEINER, F.M., SCHLICK-STEINER, B.C., SANETRA, M., LJUBOMIROV, T., ANTONOVA, V., CHRISTIAN, E. & STAUFFER, C. 2005: Towards DNA-aided biogeography: An example from *Tetramorium* ants (Hymenoptera, Formicidae). – Annales Zoologici Fennici 42: 23-35.

YUSUPOV, Z.M. 2009: An ant species of the genus *Pyramica* (Hymenoptera, Formicidae) new for the fauna of Russia and the Northern Caucasus. – Entomological Review 89: 123-124.