15(7).	With head in full-face view, anterior margin of scape with projecting curved hairs, of which one or more, distal to subbasal bend, distinctly curve toward base of scape. These hairs may be simple, spatulate, spoon-shaped, or wire-like	16
_	With head in full-face view, anterior margin of scape without projecting hairs that distinctly curve toward base of scape. Scape edge may have elongate simple	
(- -)	straight or flagellate projecting hairs present; or may have entirely anteriorly or apically directed short hairs; or lacking hairs	21
16(15).	Pronotal humeral hair present, may be filiform, flagellate, remiform, or clavate; humeral hair always distinctly differentiated from any other pilosity that may be present on dorsal pronotum	17
_	Pronotal humeral hair absent; humerus without a hair that is distinctly differentiated	20
17(16).	from any other pilosity that may be present on dorsal pronotum	20
	spongiform cuticular carina, but if weakly spongiform strip occurs then its maximum depth only fraction of depth of peduncle. Disc of postpetiole usually sculptured at least in part, only rarely mostly smooth	18
-	Ventral surface of petiole in profile with deep, conspicuous and very obviously spongiform curtain, its maximum depth at least half that of peduncle and usually more. Disc of	10
	postpetiole completely unsculptured and glassy smooth Pyramica alb	erti
18(17).	Metapleuron and side of propodeum entirely reticulate-punctate	19
- ` ´	Metapleuron and side of propleuron mostly or entirely smooth and shinning	
	Pyramica cincinn	ata
19(18).	Pronotal humeral hair elongate and freely projecting, slightly flattened apically and more or less straight. Scape in dorsal view slender, broadest point distinctly distal	
	of midlenght. Anterior margin of clypeus very shallowly convex in full-face view.	hia
	Disc of postpetiole not entirely densely reticulate-punctate Pyramica urrho Pronotal humeral hair very short, clavate. Scape in dorsal view broad and flattened,	oiu
_	broadest point proximal of midlenght, at or just distal of subbasal bend. Anterior margin of clypeus transverse to very shallowly concave in full-face view. Disc of	
	postpetiole entirely densely reticulate-punctate Pyramica me	trix
20(16).	Promesonotum, side of mesosoma, and disc of postpetiole finely reticulate-punctate. Head in profile incredible dorsoventrally flattened; at eye level depth of head	
	capsule scarcely more than twice vertical diameter of the eye. Ventral margin of petiole lacking curtain of lamellate or spongiform tissue Pyramica depression	ceps
-	Promesonotum and side of mesosoma smooth and shining, disco of pospetiole not reticulate-punctate. Head in profile not strongly dorsoventrally flattened; at eye	•
	level depth of head capsule distinctly more than twice vertical diameter of eye. Ventral margin of petiole with lamellate curtain that extends entire length of	
	segment Pyramica thax	cteri
21(15).	With head in full-face view dorsolateral margin behind level of eye with laterally projecting	
(,	hairs present; at least an apicoscrobal hair but often more along margin	22
	With head in full-face view dorsolateral margin behind level of eye without laterally	
	projecting hairs of any form; any hairs that do occur are minute and closely appressed, not at all projecting	24
22(21)		
22(21).	extends entire length of sclerite. With postpetiole in profile ventral spongiform lobe	
	either completely absent or reduced to minute triangular vestige anteriorly on	tata
	sternite	ıuıu
-	Midline of clypeal dorsum not raised into a high longitudinal crest that extends length of sclerite. With postpetiole in profile ventral spongiform lobe fully developed, basally extending length of sternite and conspicuously convex apically	23
	basany extending length of sterrine and conspicuously convex apicany	