

- Head distinctly lighter than mesosoma (exceptions in *O. infandus* only occur, if both parts are very dark brown). Petiole less stout. (Figs. 3, 5, 9, 11, 21, 22) **9**
- 9 Gaster tergites 1 and 2 partly yellowish brown, tergite 1 anteriorly low, strongly flattened; impression, if pit-shaped, at or behind mid-length. Petiole with very elongate spine. Striation of pronotum variable (transverse or longitudinal). Species from the central and eastern Philippines (absent from Luzon and Mindoro). (Figs. 5, 12, 17) ***Odontomachus alius* sp.n.**
- Gaster tergites 1 and 2 dark (blackish) brown, tergite 1 anteriorly comparatively high; impression, if pit-shaped, before mid-length (Fig. 22). Petiolar spine shorter (Figs. 9, 22). Striation of pronotum longitudinal, usually in loops (Figs. 15, 23). **10**
- 10 Striation of head almost complete. Head usually light to dark brown. Petiolar spine curved. Species widely distributed on Luzon and Mindoro. (Figs. 3, 9, 15) ***Odontomachus infandus***
- Striation of head posteriorly reduced. Head yellowish orange, strongly contrasting with mesosoma. Petiolar spine straight. Species from southern Luzon. (Figs. 21 - 23) ***Odontomachus* sp. 1**



Figs. 21 - 23: *Odontomachus* sp. 1: (21) head, full face view; (22, below) habitus, lateral view; (23, below) habitus, dorsal view. Scales = 1 mm.

