

*strigatus* group is restricted to 20° and 30° south (Sanhudo et al. 2007). The *strigatus* group also lacks species with wide distributions (Mayhé-Nunes and Jaffé 1998) as is found in the *rimosus* complex.

Apparently no new species of the *strigatus* group have been described since Kempf's revision (1964), although the recently described *C. muelleri* shows similarities to the *strigatus* species complex, and along with *C. longiscapus*, *C. costatus* and *C. wheeleri* may be related (Schultz et al. 2002). A similar new genus *Mycetagroicus*, with three new species has been recently described (Brandão and Mayhé-Nunes 2001).

### GLOSSARY

**Anteclypeus** (= "apron"): the anterior portion of the clypeus attached to the labrum. **Dorsopropodeum**: the dorsal surface of the propodeum.

**Frontovertexal corner**: the posterolateral angle between frons and vertex.

**Notopropodeal groove**: (= "metanotal groove") a transverse groove on the noto-propodeal fusion.

**Notopropodeal fusion**: In workers, the tergal fusion of the thoracic notum and the propodeum.

**Opisthogaster** (adj. opisthogastral): (= "gaster") Abd IV to pygidium.

**Paraclypeal teeth**: (= "parafrontal teeth" - Kempf 1964, 1965), referring to the anterior teeth-like processes on the clypeus.

### ACKNOWLEDGMENTS

We would like to thank Roy Snelling and Weiping Xie for the loan of specimens of *Cyphomyrmex faunulus*. Two anonymous reviewers provided important comments. The research was supported by a grant from the National Geographic Society to Robert Anderson and the Ernst Mayr Fund of the Museum of Comparative Zoology.

### LITERATURE CITED

- Bolton, B., G. Alpert, P. Ward, and P. Naskrecki. 2007. *Bolton's Catalog of Ants of the World: 1758–2005*. Harvard University Press, available on CD.
- Brandão, C. and A. Mayhé-Nunes. 2001. A new fungus-growing ant genus, *Mycetagroicus* gen. With the description of three new species and comments on the monophyly of the Attini (Hymenoptera: Formicidae). *Sociobiology* 38: 639–665.
- Fernández, F. and E. Palacio. 1995. Hormigas de Colombia IV: Nuevos registros de géneros y especies. *Caldasia* 17: 587–596.
- Kempf, W. W. 1964. A revision of the Neotropical ants of the genus *Cyphomyrmex* Mayr. Part I. Group of *strigatus* Mayr (Hym. Formicidae). *Studia Entomologica* 7: 1–44.
- . 1965. A revision of the Neotropical fungus-growing ants of the genus *Cyphomyrmex* Mayr. Part II: group of *rimosus* (Spinola) (Hym. Formicidae). *Studia Entomologica* 8: 161–200.
- Mayhé-Nunes, A. and K. Jaffé. 1998. On the biogeography of Attini (Hymenoptera: Formicidae). *Ecotropicos* 11: 45–54.
- Sanhudo, C., A. Mayhé-Nunes, and C. Brandão. 2007. Quem são as *Cyphomyrmex* (Myrmicinae: Attini). *Biológico* 69: 433–434.
- Schultz, T. and S. Brady. 2008. Major evolutionary transitions in ant agriculture. *Proceedings of the National Academy of Sciences* 105: 5435–5440.
- , S. Solomon, U. Mueller, P. Villesen, J. Boomsma, R. Adams, and B. Norden. 2002. Cryptic speciation in the fungus-growing ants *Cyphomyrmex longiscapus* Weber and *Cyphomyrmex muelleri* Schultz and Solomon, new species (Formicidae, Attini). *Insectes Sociaux* 49: 331–343.
- Serna, F. and W. Mackay. 2010. A descriptive morphology of the ant genus *Procryptocerus* (Hymenoptera: Formicidae). *Journal of Insect Science* (in press).
- Snelling, R. R. and J. T. Longino. 1992. Revisionary notes on the fungus-growing ants of the genus *Cyphomyrmex*, *rimosus* group (Hymenoptera: Formicidae: Attini). Pp. 479–494 in: Quintero, D., and A. Aiello eds. *Insects of Panama and Mesoamerica: selected studies*. Oxford University Press, Oxford. 692 pp.