

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank Seán Brady and Robert Kallal for comments on earlier drafts of this manuscript. JSL was supported by the National Science Foundation under grant DEB-0743542; GMD was supported by the Russian Foundation for Basic Research under grant 11-04-00421; VP was supported by the French National Institute for Universe Sciences under grant Interrvie NOVAMBRE.

LITERATURE CITED

1. Archibald SB, Cover SP, Moreau CS. 2006. Bulldog ants of the Eocene Okanagan Highlands and history of the subfamily (Hymenoptera: Formicidae: Myrmeciinae). *Ann. Entomol. Soc. Am.* 99:487–523
2. Archibald SB, Farrell BD. 2003. Wheeler's dilemma: Proceedings of the Second Paleontological Congress. *Acta Zool. Cracov.* 46(Suppl.-Fossil Insects):17–23
3. Archibald SB, Johnson KR, Mathewes RW, Greenwood DR. 2011. Intercontinental dispersal of giant thermophilic ants across the Arctic during early Eocene hyperthermals. *Proc. R. Soc. B* 278:3679–86
4. Aria C, Perrichot V, Nel A. 2011. Fossil Ponerinae (Hymenoptera: Formicidae) in Early Eocene amber of France. *Zootaxa* 2870:53–62
5. Baroni Urbani C, Wilson EO. 1987. The fossil members of the ant tribe Leptomyrmecini (Hymenoptera: Formicidae). *Psyche* 94:1–8
6. Bolton B. 2003. Synopsis and classification of Formicidae. *Mem. Am. Entomol. Inst.* 71:1–370
7. Bolton B. 2012. *AntCat: An Online Catalog of the Ants of the World*. <http://www.antcat.org>
8. Brandão CR, Martins-Neto R, Vulcano VA. 1989. The earliest known fossil ant (first Southern Hemisphere Mesozoic record) (Hymenoptera: Formicidae: Myrmeciinae). *Psyche* 96:195–208
9. Brady SG, Schultz TR, Fisher BL, Ward PS. 2006. Evaluating alternative hypotheses for the early evolution and diversification of ants. *Proc. Natl. Acad. Sci. USA* 103:18172–77
10. Bromley RG, Buatois LA, Genise JF, Labandeira CC, Mángano MG, et al. 2007. Comments on the paper “Reconnaissance of Upper Jurassic Morrison Formation ichnofossils, Rocky Mountain Region, USA: paleoenvironmental, stratigraphic, and paleoclimatic significance of terrestrial and freshwater ichnocoenoses” by Stephen T. Hasiotis. *Sediment. Geol.* 200:141–50
11. Brown WL Jr, Carpenter FM. 1978. A restudy of two ants from the Sicilian amber. *Psyche* 85:417–23
12. Carpenter FM. 1929. A fossil ant from the Lower Eocene (Wilcox) of Tennessee. *J. Wash. Acad. Sci.* 19:300–1
13. Carpenter FM. 1930. The fossil ants of North America. *Bull. Mus. Comp. Zool. Harv.* 70:1–66
14. Clementz M, Bajpai S, Ravikant V, Thewissen JGM, Saravanan N, et al. 2011. Early Eocene warming events and the timing of terrestrial faunal exchange between India and Asia. *Geology* 39:15–18
15. Cockerell TDA. 1915. British fossil insects. *Proc. U.S. Natl. Mus.* 49:469–99
16. Crozier RH, Jermini LS, Chiotis M. 1997. Molecular evidence for a Jurassic origin of ants. *Naturwissenschaften* 84:22–23
17. Dierick M, Cnudde V, Masschaele B, Vlassenbroeck J, Van Hoorebeke L, Jacobs P. 2007. Micro-CT of fossils preserved in amber. *Nucl. Instrum. Methods Phys. Res. A* 580:641–43
18. Dlussky GM. 1975. Superfamily Formicoidea. See Ref. 88, pp. 114–22
19. Dlussky GM. 1983. A new family of Upper Cretaceous Hymenoptera: an “intermediate link” between ants and Scolioidea. *Paleontol. J.* 17:63–76
20. Dlussky GM. 1987. New Formicoidea (Hymenoptera) of the Upper Cretaceous. *Paleontol. J.* 20:146–50
21. Dlussky GM. 1988. Ants from (Paleocene?) Sakhalin amber. *Paleontol. J.* 22:50–61

19. First description of the Armaniidae.
