



Figs. 21 – 23. *Dilobocondyla fouqueti*, Worker: (21) Head in full-face view; (22) Body, lateral view; (23) Body, dorsal view.

ETYMOLOGY. The species is named in honour of Professor Seiki Yamane, who generously donated it for present study.

REMARKS. In *D. borneensis*, *D. didita*, *D. eguchii*, *D. gaoyureni*, *D. karnyi*, *D. propotriangulatus* and *D. sebesiana* too the frontal carinae do not continue to the posterior corners of the head, but *Dilobocondyla yamanei* sp. nov. can be separated from these by the following combination of

Figs. 24 – 26. *Dilobocondyla fouqueti*, Gyne: (24) Head in full-face view; (25) Body, lateral view; (26) Body, dorsal view.

features: ferruginous body colour; shorter petiole; opaque body; presence of 9 rugae between frontal carinae; and also by body length (TL 4.83 mm). The most similar taxon to this species is *D. karnyi*, from which it can be easily distinguished on the basis of the following combination of characters: head broader than long (longer than broad in *D. karnyi*), length 4.83 mm (4.5 mm in *D. karnyi*), and petiole less than twice as long as broad (more than twice as long as broad in *D. karnyi*). Other significant differences which separate it from *D. karnyi* include: clypeus with a strong median and two lateral carinae, its anterior border strongly emarginate in the middle; propodeum with short basal part which is almost equal to declivity