

Lasius piliferus Seifert, 1992, Sant Julià de Lòria: Coll de Jou, carretera de Fontaneda (42°27.50'N; 1°29.00'E - 1100 m a.s.l), 02.VIII.2008, leg. A. Bernadou, det. X. Espadaler, workers collected.

Tapinoma madeirense Forel, 1895, Sant Julià de Lòria: Coll de Jou, carretera de Fontaneda (42°27.50'N; 1°29.00'E - 1100 m a.s.l), 02.VIII.2008, leg. A. Bernadou, det. X. Espadaler, workers collected.

Temnothorax lichtensteini (Bondroit, 1918), Sant Julià de Lòria: Coll de Jou, carretera de Fontaneda (42°27.50'N; 1°29.00'E - 1100 m a.s.l), 02.VIII.2008, leg. A. Bernadou, det. X. Espadaler, workers collected.

Temnothorax niger (Forel, 1894), Sant Julià de Lòria: Coll de Jou, carretera de Fontaneda (42°27.50'N; 1°29.00'E - 1100 m a.s.l), 02.VIII.2008, leg. A. Bernadou, det. X. Espadaler, workers collected.

Temnothorax nigriceps (Mayr, 1855), Sant Julià de Lòria: Coll de Jou, carretera de Fontaneda (42°27.50'N; 1°29.00'E - 1100 m a.s.l), 02.VIII.2008, leg. A. Bernadou, det. X. Espadaler, workers collected.

Table 1. Checklist of the ant species of Andorra. The list is arranged alphabetically by subfamily, genus and species. Species names in bold characters refer to species recorded for the first time in Andorra. Bibliographic references are as follows: a = Bernadou et al. 2006a; b = Bernadou et al. 2006b; c = Espadaler et al. 2008 *Hypoponera eduardi* was misidentified with *Hypoponera punctatissima* in this study (Espadaler, pers. com.); d = Bernadou et al. 2010; e = Bernadou 2009; f = Seifert 1992; g = Santschi 1919; h = Espadaler 1997; i = Collingwood and Yarrow 1969; j = Leppänen et al. 2011; k = Cournault and Aron 2009; l = Bagherian et al. 2012; m = Rösler 1951.

Subfamilies	Scientific valid name		References
DOLICHODERINAE	<i>Dolichoderus quadripunctatus</i>	(Linnaeus, 1771)	c
	<i>Tapinoma erraticum</i>	(Latreille, 1798)	a, b, c, e, k
	<i>Tapinoma madeirense</i>	Forel, 1895	
FORMICINAE	<i>Camponotus aethiops</i>	(Latreille, 1798)	c
	<i>Camponotus cruentatus</i>	(Latreille, 1802)	c
	<i>Camponotus herculeanus</i>	(Linnaeus, 1758)	a, b, e
	<i>Camponotus lateralis</i>	(Olivier, 1792)	
	<i>Camponotus ligniperda</i>	(Latreille, 1802)	a, b, c, e, i
	<i>Camponotus piceus</i>	(Leach, 1825)	
	<i>Camponotus truncatus</i>	(Spinola, 1808)	c
	<i>Formica decipiens</i>	Bondroit, 1918	a, b, e
	<i>Formica exsecta</i>	Nylander, 1846	
	<i>Formica foreli</i>	Bondroit, 1918	a, b, e
	<i>Formica frontalis</i>	Santschi, 1919	a, b, e
	<i>Formica fusca</i>	Linnaeus, 1758	a, b, c, e, g
	<i>Formica genardi</i>	Bondroit, 1917	c
	<i>Formica lemani</i>	Bondroit, 1917	a, b, e
	<i>Formica lugubris</i>	Zetterstedt, 1838	a, b, e, i
<i>Formica picea</i>	Nylander, 1846	a, b, e	
<i>Formica pratensis</i>	Retzius, 1783	a, b, e	