

Subfamilies	Scientific valid name	References
	<i>Formica pressilabris</i>	a, b, e
	<i>Formica rufa</i>	a, b, e, g
	<i>Formica rufibarbis</i>	a, b, c, e, i
	<i>Formica sanguinea</i>	a, b, e
	<i>Lasius alienus</i>	a, b, c, e
	<i>Lasius brunneus</i>	c, e
	<i>Lasius distinguendus</i>	c, h
	<i>Lasius flavus</i>	(Fabricius, 1782)
	<i>Lasius fuliginosus</i>	(Latreille, 1798)
	<i>Lasius grandis</i>	Forel, 1909
	<i>Lasius mixtus</i>	(Nylander, 1846)
	<i>Lasius niger</i>	(Linnaeus, 1758)
	<i>Lasius paralienus</i>	Seifert, 1992
	<b><i>Lasius piliferus</i></b>	Seifert, 1992
	<i>Lasius platythorax</i>	Seifert, 1991
	<i>Plagiolepis pygmaea</i>	(Latreille, 1798)
	<i>Plagiolepis xene</i>	Stärcke, 1936
MYRMICINAE	<b><i>Aphaenogaster gibbosa</i></b>	(Latreille, 1798)
	<i>Aphaenogaster subterranea</i>	(Latreille, 1798)
	<i>Crematogaster scutellaris</i>	(Olivier, 1791)
	<i>Leptothorax acervorum</i>	(Fabricius, 1793)
	<i>Leptothorax muscorum</i>	(Nylander, 1846)
	<i>Messor structor</i>	(Latreille, 1798)
	<i>Myrmecina graminicola</i>	(Latreille, 1802)
	<i>Myrmica lobulicornis</i>	Nylander, 1857
	<i>Myrmica rubra</i>	(Linnaeus, 1758)
	<i>Myrmica ruginodis</i>	Nylander, 1846
	<i>Myrmica sabuleti</i>	Meinert, 1861
	<i>Myrmica scabrinodis</i>	Nylander, 1846
	<i>Myrmica schencki</i>	Emery, 1895
	<i>Myrmica specioides</i>	Bondroit, 1918
	<i>Myrmica spinosior</i>	Santschi, 1931
	<i>Myrmica sulcinodis</i>	Nylander, 1846
	<i>Myrmica wesmaeli</i>	Bondroit, 1918
	<i>Pheidole pallidula</i>	(Nylander, 1849)
	<i>Pyramica tenuipilis</i>	(Emery, 1915)
	<i>Solenopsis</i> sp.	Westwood, 1840
	<i>Stenamma striatulum</i>	Emery, 1895
	<i>Strongylognathus testaceus</i>	(Schenck, 1852)
	<i>Temnothorax affinis</i>	(Mayr, 1855)
	<i>Temnothorax gredosi</i>	(Espadaler & Collingwood, 1982)
	<i>Temnothorax kraussei</i>	(Emery, 1916)
	<b><i>Temnothorax lichtensteini</i></b>	(Bondroit, 1918)
	<i>Temnothorax nadigi</i>	(Kutter, 1925)