

- 4 Distal 2/3 of basal margin of mandible straight, proximal 1/3 narrowed toward base of mandible; anterior clypeal margin straight; petiole clearly smaller than postpetiole (Fig. 14C) (Lombok) ***A. minipetiolus* sp. n.**
- Basal margin of mandible feebly concave; anterior clypeal margin concave; petiole larger than or as large as postpetiole (Figs 2C, 20C) **5**
- 5 Promesonotum strongly convex and sloping gradually to metanotal groove; subpetiolar process with angulate posteroventral corner (Fig. 2C); antennal scape relatively long (SI 78–86) (Bali)..... ***A. baliensis* sp. n.**
- Mesosoma almost flat dorsally or feebly convex; subpetiolar process lower, with its posteroventral corner rounded (Fig. 20C); antennal scape relatively short (SI 57–78) (Vietnam and Thailand)..... ***A. wiwatwitayai* sp. n.**
- 6 Vertex with 2 long standing hairs mixed with a few short appressed hairs; promesonotum with few appressed hairs and a few decumbent hairs (Fig. 1C) (Sabah) ***A. appressipilosus* sp. n.**
- Vertex and promesonotum with dense standing or decumbent hairs (Figs 10C, 13C, 15C, 16) **7**
- 7 Promesonotum entirely sculptured (punctate or reticulate) **8**
- Promesonotum largely smooth and shiny..... **9**
- 8 Promesonotum sparsely reticulate and shiny or superficially micropunctate (entire promesonotum looks shiny); antennal scape superficially shagreened; petiole cylindrical, clearly longer than high (Fig. 5C) (S. Thailand)..... ***A. cylindripetiolus* sp. n.**
- Promesonotum densely punctate and matt; antennal scape micropunctate; petiole round or subangulate, almost as long as high (Fig. 17D) (Vietnam and N. Thailand) ***A. thailandianus* Terayama & Kubota**
- 9 With head seen in profile and dorsal view occipital corner produced as a small lobe (Fig. 9C) (Sulawesi) ***A. goniocippus* sp. n.**
- With head seen in profile occipital corner without such a lobe (Figs 15C, 23C) **10**
- 10 Masticatory margin of mandible with large acute apical tooth followed by a series of 6–7 denticles of two sizes, the larger alternating with 1–2 smaller (Fig. 11C); gap between anterior clypeal margin and mandibles relatively small or indistinct, with maximum width shorter than maximum width of mandible (S. Japan and Taiwan)..... ***A. lifuiiae* Terayama**
- Masticatory margin of mandible with large acute apical tooth followed by a medium-sized subapical tooth, 2–5 denticles, and a medium-sized basal tooth (Fig. 17B); gap between anterior clypeal margin and mandibles relatively large and distinct, with maximum width longer than maximum width of mandible..... **11**
- 11 Subpetiolar process subtriangular, its apex directed downward, anteroventrally not angulate (Fig. 21C); postpetiole elevated posteriorly; its posterior face much steeper than anterior face (Fig. 21C) (Sumatra) ***A. itoi* sp. n.**