

- Subpetiolar process subrectangular or low, convex in its ventral outline, anteroventrally always angulate (Figs 1C, 9C, 23C); postpetiole with roundly convex dorsal outline (Figs 9C, 23C) **12**
- 12 Subpetiolar process rectangular, ventrally with anterior and posterior corners (Figs 6C, 23C) **13**
- Subpetiolar process low, its ventral outline convex, almost straight or feebly concave, and anteroventral corner acutely angulate (Figs 15C, 19C) **14**
- 13 Posteroventral corner of subpetiolar process bluntly angulate (not spiniform) (Fig. 6C); dorsal outline of propodeum weakly convex; metapleural gland bulla weakly sculptured and shiny (Vietnam) ***A. eguchii* sp. n.**
- Posteroventral corner of subpetiolar process acutely produced below (spiniform) (Fig. 23C); dorsal outline of propodeum straight; metapleural gland bulla strongly sculptured (puncto-reticulate) and opaque (Borneo and Java).
..... ***A. jawadwipa* sp. n.**
- 14 Promesonotum in profile weakly convex (Fig. 19C); propodeal dorsum feebly convex; petiole sessile with low node, its posterior face encircled with a thin carina; postpetiole almost as long as petiole (Thailand) ***A. wilaiae* sp. n.**
- Promesonotum in profile strongly convex and forming a dome (Fig. 15C); propodeal dorsum clearly straight; petiole subsessile with high node, its posterior face not encircled with a carina; postpetiole slightly shorter than petiole (Philippines) ***A. pilosus* sp. n.**
- 15 Mandible with 3 teeth including apical and basal teeth (Fig. 18B). **16**
- Mandible with 4 teeth including apical and basal teeth (Fig. 13D). **17**
- 16 With head seen in profile occipital corner round; promesonotum in profile strongly convex and forming a dome (Fig. 18D); petiole almost as long as high, with high node, and almost as long as postpetiole (Vietnam and Thailand) ***A. watanasiti* sp. n.**
- With head seen in profile occipital corner roundly angulated (almost right-angled) (Figs 4C); promesonotum in profile with weakly convex or almost flat dorsal outline (Fig. 4C); petiole clearly longer than high, with low node, and slightly longer than postpetiole (Vietnam and Thailand) ... ***A. concavus* sp. n.**
- 17 Dorsal and lateral faces of pronotum largely smooth and shiny; petiole almost as long as high; head slightly longer than or almost as long as broad (CI 90-100) **18**
- Dorsal and lateral faces of pronotum entirely sculptured (superficially puncto-reticulate, punctate to reticulate); petiole cylindrical, distinctly longer than high; head slightly shorter than broad (CI 102-112) **20**
- 18 Anterior clypeal margin distinctly concave; metanotal groove distinct, deep; foretibia relatively short, its length less than 0.5 times of head width; mesopleuron finely punctate without longitudinal rugulae (N. Vietnam)
..... ***A. brevipodus* sp. n.**