



**Figure 15.** *Aenictus pilosus* (holotype). **A** Head in full-face view **B** body in dorsal view **C** body in profile.

junction angulated, nearly right-angled; declivity of propodeum shallowly concave, and encircled with a rim; metapleural gland bulla relatively large, its maximum diameter about 2.5 times as long as distance between propodeal spiracle and metapleural gland bulla. Petiole slightly longer than high, elevated posteriorly; subpetiolar process generally very low, with its anteroventral corner acutely angulate, and ventral margin feebly concave. Postpetiole slightly smaller than petiole, with its dorsal outline convex.

Head and gaster smooth and shiny; basal half of antennal scape densely microreticulate but apical half smooth and shiny; mandible finely striate. Promesonotum smooth and shiny except for anteriormost portion punctate; mesopleuron with slightly irregular rugae; metapleuron, and propodeum wrinkled and reticulate. Petiole entirely reticulate; postpetiole superficially reticulate and shiny. Basal 1/3 of femora sculptured, apical 2/3 (swollen area) smooth and shiny; tibiae microreticulate, partly smooth and shiny.

Head and mesosoma dorsally with relatively dense standing hairs; longest pronotal hair 0.33–0.35 mm long; legs with dense long decumbent hairs. Head reddish brown;