

- Triton palustris*. (Warty Eft.)
 — *punctatus*. (Common Eft.)

PISCES.

- Perca fluviatilis*. (Perch.)
Cottus Gobio. (Bullhead.)
Cyprinus Carpio. (Common Carp.)
 — *Gobio*. (Gudgeon.)
 — *Tinca*. (Tench.)
 — *Rutilus*. (Roach.)
 — *Leuciscus*. (Dace.)
 — *Cephalus*. (Chub.)
 — *Phoxinus*. (Minnow.)
Cobitis barbatula. (Bearded Loach.)
Esox Lucius. (Pike.)
Salmo Salar. (Common Salmon.)
 — *Fario*. (Common Trout.)
Thymallus vulgaris. (Grayling.)
Platessa Flesus. (Flounder.) A single specimen caught with rod
 and line Dec. 1839, in river Lug, below Mordiford Bridge.
Anguilla acutirostris. (Sharp-nosed Eel.) } In river Lug.
 — *latirostris*. (Broad-nosed Eel.) }

XXII.—*Monograph of the Dorylidæ, a Family of the Hymenoptera Heterogyna*. By W. E. SHUCKARD, Esq.

THE discovery of an insect that will, I expect, help to clear up the difficulty which has hitherto attended the completion of these genera, as yet consisting of males only, has induced me to undertake the present monograph. Although the materials with which I entered upon this task were rather scanty, they have grown upon my hands and are now coextensive with the metropolitan collections; and when we know that these comprise the collections of many individuals, all much attached to the order Hymenoptera, we must conclude that these genera are naturally poor in individuals, although the number of species that I produce far exceed all that have been hitherto described. In the genus *Dorylus* three species only have yet been noticed, two African and one Indian, but it is very questionable if one of the African species may not, understood under the name of *D. helvolus*, consist of many species from that quarter of the globe which constitute the majority of the family, for Africa is evidently