

spinosis; thorace spinis duabus acutis elongatis armato; abdomine cordato.

*Worker.* Length  $2\frac{1}{2}$  lines. Opaque-black; the head and thorax above, with a coarse irregular striation, intermixed with a rough granulation, the margins of the head crenulated, the posterior angles acute and slightly produced. Thorax armed posteriorly with two stout diverging spines. Abdomen oblong-cordate, finely and irregularly striated; sprinkled with distant short white erect setæ; the nodes of the peduncle rugose.

*Hab.* Celebes (Tondano).

#### Gen. ECHINOPLA, *Smith.*

1. *Echinopla striata*, *Smith, Proc. Linn. Soc. ii. 80. 3.*

*Hab.* Celebes; Malacca.

2. *Echinopla pallipes*, *Smith, Proc. Linn. Soc. ii. 80. 2.*

*Hab.* Celebes; Borneo.

3. *ECHINOPLA DUBITATA.* *E. nigra*; capite thoraceque rugosis; abdomine ovato lævi nitido; squama in utroque latere spina horizontali; femoribus pallide testaceis.

*Worker.* Length 2 lines. Black, and thinly covered with erect black hairs; the head and thorax rather finely rugose; the antennæ pubescent, with the extreme tip pale testaceous; the eyes round and very prominent. The anterior margin of the prothorax arched with a short acute spine at the lateral angles; the thorax is deeply constricted in the middle, the metathorax rounded behind; the roughness of the thorax gives its margins a crenulated appearance; the coxæ, trochanters, and base of the femora pale rufo-testaceous; the claw-joint of the tarsi rufo-testaceous. Abdomen globose, shining, and very finely punctured.

*Hab.* Celebes (Tondano).

#### Fam. MUTILLIDÆ, *Leach.*

##### Gen. MUTILLA.

1. *Mutilla Merops*, *Smith, Supp. Journ. Proc. Linn. Soc. v. 115. 2.*

*Hab.* Gilolo; Bachian.

2. *Mutilla anthylla*, *Smith, Supp. Journ. Proc. Linn. Soc. v. 115. 4.*

*Hab.* Gilolo; Bachian.

3. *Mutilla Ianthea*, *Smith, Proc. Linn. Soc. Supp. v. 115. 3.*

*Hab.* Amboyna; Bachian.

##### Gen. METHOCA, *Latr.*

1. *METHOCA THORACICA.* *M. rufo-ferruginea*; capite abdominisque fasciis tribus nigris.