

also appears more vertical, and the general body color more infuscated or sordid.

DEALATE FEMALE. Length 2.01 mm.

Larger than the worker. Total body length obtained in the same manner as with the worker and comprising the following parts, head 0.50 mm., thorax 0.66 mm., gaster 0.85 mm. Differing from the worker largely in the usual female characters such as the possession of ocelli, compound eyes, and thoracic modifications. Other differences are mandible edentate except for the single apical tooth; eye rather large, oval, approximately 0.13 mm. at its greatest width and bearing in this width 10-12 ommatidia, placed approximately 0.05 mm. from the base of the mandible; anterior ocellus borne less than 0.10 mm. behind a transverse line connecting the posterior border of each eye; compound eyes and borders around the ocelli black; pronotal humeri more angular; epinotal declivity apparently more concave. Color similar to that of the two Georgia workers but more sordid than that of the worker from North Carolina.

The redescription of the worker is based largely on the individual from North Carolina which has been carefully compared with the type. The redescription of the female is from the single, dealated Georgia individual.

Type locality — North America (very probably collected in one of the Carolinas by Dr. Christian Zimmerman).

Other localities — 3 miles from Holly Springs (Wake County), North Carolina, Merle W. Wing and Savannah, Georgia, H. T. Vanderbilt.