

and fainter laterad; those of disc about equal to those of occiput; punctures of upper mesopleural plate slightly finer than those of adjacent portion of pronotum, separated by slightly less than a puncture diameter; punctures of lower half of mesopleura, entire metapleura, lateral and posterior faces of epinotum obscure, scattered, the integument opaque, minutely granulose; punctures of scutellum slightly elongate, equal to those of mesoscutum, separated by a puncture diameter or less, the integument slightly more shiny than elsewhere; punctures of basal face of epinotum coarse, deep, subcontiguous.

Wings fuscous infusate, except for clear submarginal cell; marginal cell closed and appendiculate apically, about 3.5 times as long as wide; submarginal cell about three-fourths as long as marginal; otherwise as figured by Kempf (1959:93) for the male.

Described from a single alate female, associated with forty workers, from Pucallpa, Loreto, Peru, alt. 200 m., collected November 11-20, 1964 by J. Schunke. All specimens in the collection of the Los Angeles County Museum.

LITERATURE CITED

KEMPF, W. W.

1951. A taxonomic study on the ant tribe Cephalotini. *Revista de Entomologia*, 22:1-244.

1959. Sobre algumas formigas "Cephalotini" do Museu de Oxford. *Revista Brasileira Biologia*, 19:91-98.