

with this smaller size is a relatively longer antennal scape and a slightly duller integument (especially on the head). The western populations are highly plastic (perhaps in response to greater variation in habitats and climate) and individuals within a single colony sample may possess either a long or short scape, or one of intermediate length.

## REFERENCES

- Brown, W. L., Jr.** 1950. The status of two common North American carpenter ants. *Ent. News* 61:157-161.
- Creighton, W. S.** 1950. The ants of North America. *Bull. Mus. Comp. Zool.* 104:1-585.
- . 1952. Studies on Arizona ants (4). *Camponotus (Colobopsis) papago*, a new species from southern Arizona. *Psyche* 59:148-162.
- . 1965. Studies on southwestern ants belonging to *Camponotus*, subgenus *Myrmobrachys*. *Amer. Mus. Novit.* 2239:1-9.
- and **R. R. Snelling.** 1966. The rediscovery of *Camponotus yogi* Wheeler. *Psyche* 73:187-195.
- Gregg, R. E.** 1963. The ants of Colorado. Univ. Colo. Press, Denver, xvi + 792 pp.
- Roger, J.** 1863. Die neu aufgeführten Gattungen und Arten meins Formiciden-Verzeichnisses. *Berlin Ent. Zeit.* 7:131-214.
- Smith, M. R.** 1953. A new *Camponotus* in California apparently inhabiting live oak, *Quercus* sp. *Jour. N. Y. Ent. Soc.* 61:211-214.
- Wheeler, W. M.** 1910a. The North American forms of *Camponotus fallax* Nylander. *Jour. N. Y. Ent. Soc.* 18:216-232.
- . 1910b. The North American ants of the genus *Camponotus* Mayr. *Ann. N. Y. Acad. Sci.* 20:295-354.