

and ocelli, roughened, with scattered, obscure punctures of variable size
*placodops* Forel

Mesoscutal disc with entire median area impunctate; frontal lobes somewhat
 shining, with fine, dense, evenly spaced punctures; vertex between eyes and
 ocelli with fine, distinct, close punctures.....*melliger* Forel

Type material of the species discussed above has been made available through the courtesy of the following: C. Besuchet and C. Ferriere, Muséum d'Historie Naturelle, Geneva; W. L. Brown, Jr., Museum of Comparative Zoology; J. G. Rozen, Jr., American Museum of Natural History; D. R. Smith, United States National Museum. To each of these gentlemen, I offer my most profound thanks for their cooperation. The figures were prepared by Ruth Ann DeNicola under assistance rendered by Grant No. 4494, the Penrose Fund, of the American Philosophical Society.

LITERATURE CITED

- CREIGHTON, W. S. 1950. The ants of North America. Harvard, Mus. Comp. Zool., Bull., 104:1-585.
 CREIGHTON, W. S., AND R. H. CRANDALL. 1954. New data on the habits of *Myrmecocystus melliger* Forel. C. C. N. Y., The Biol. Rev., 16:2-6.
 GREGG, R. E. 1963. The ants of Colorado. Univ. Colo. Press, 792 pp.
 WHEELER, W. M. 1908. Honey ants, with a revision of the American Myrmecocysti. Amer. Mus. Nat. Hist., Bull., 24:345-397.

Accepted for publication February 24, 1969