

seven cells from a single worker female collected in Arizona revealed $2K = 14\bar{M} + 2\bar{A}$ (Tables 1, 2). Our karyotypes were essentially identical to those published by CROZIER (1970: Fig. 1B).

Acknowledgements. — We thank Dr. OSCAR F. FRANCKE and Mr. SCOTT STOCKWELL for aid in the collection of ants. Dr. ANDRÉ FRANCOEUR kindly identified the *Leptothorax rugatulus*. Dr. M. KENT RYLANDER provided laboratory space and equipment for which we are grateful. Drs. BARRY BOLTON, ANDRÉ FRANCOEUR, JAMES TRAGER, and three reviewers for the College of Agriculture, Texas Tech University, provided useful comments and suggestions on the manuscript. This research was partially supported by the Texas Department of Agriculture and a summer research fellowship from The Graduate School, Texas Tech University and is contribution No. T-10-185 from the College of Agricultural Sciences, Texas Tech University.

REFERENCES

- BOLTON B., 1976. — *The ant tribe Tetramoriini (Hymenoptera Formicidae). Constituent genera, review of smaller genera, and revision of Triglypbothrix Forel.* Bull. Brit. Mus. (Nat. Hist.), Entomol., 34: 281-379.
- 1985. — *The ant genus Triglypbothrix Forel a synonym of Tetramorium Mayr (Hymenoptera: Formicidae).* J. Nat. Hist., 19: 243-248.
- BUSCHINGER A., 1982. — *Leptothorax jaberii n. sp., an apparently parasitic ant from Jasper National Park, Canada (Hymenoptera: Formicidae).* Psyche, 89: 197-209.
- BUSCHINGER A., FRANCOEUR A. and FISCHER K., 1980. — *Functional monogyny, sexual behavior, and karyotype of the guest ant, Leptothorax provancheri Emery (Hymenoptera, Formicidae).* Psyche, 87: 1-12.
- COKENDOLPHIER J. C. and BROWN J. D., 1985. — *Air-dry method for studying chromosomes of insects and arachnids.* Entomol. News, 96: 114-118.
- COKENDOLPHIER J. C. and FRANCKE O. F., 1985. — *Karyotype of Conomyrma flava (McCook) (Hymenoptera: Formicidae).* J. New York Entomol. Soc., 92(4): 349-351.
- CROZIER R. H., 1970. — *Pericentric rearrangement polymorphism in a North American Dolichoderine ant (Hymenoptera: Formicidae).* Can. J. Genet. Cytol., 12: 541-546.
- , 1975. — *Vol. 3: Insecta, 7. Hymenoptera.* Pp. I-V, 1-95. In: «Animal Cytogenetics» (B. John ed.), Gebrüder Berlin, Borntraeger, Stuttgart.
- , 1977. — *Genetic differentiation between populations of the ant Apbaenogaster «rudis» in the southeastern United States.* Genetica, 47: 17-36.
- FRANCOEUR A., 1986. — *Deux nouvelles fourmis Néarctiques: Leptothorax retractus et L. sphagnicolus (Formicidae, Hymenoptera).* Can. Entomol., 118: 1151-1164.
- FRANCOEUR A., LOISELLE R. and BUSCHINGER A., 1985. — *Biosystematique de la tribu Leptothoracini (Formicidae, Hymenoptera). I. Le genre Formicocoxenus de la région boréarctique.* Naturaliste Can., 112: 343-403.
- GLANCEY B. M., ROMAÏN M. K. ST. and CROZIER R. H., 1976. — *Chromosome numbers of the red and the black imported fire ants, Solenopsis invicta and S. richteri.* Ann. Entomol. Soc. Am., 69: 469-470.
- GOÑI B., ZOLESSI L. C. DE and IMAI H. T., 1983. — *Karyotypes of thirteen ant species from Uruguay (Hymenoptera, Formicidae).* Caryologia, 36: 363-371.
- HAUSCHTECK-JUNGEN E. and JUNGEN H., 1983. — *Ant chromosomes II. Karyotypes of western Palearctic species.* Insect Soc., 30: 149-169.
- HUNG A. C.-F., IMAI H. T. and KUBOTA M., 1972. — *The chromosomes of nine ant species (Hymenoptera: Formicidae) from Taiwan, Republic of China.* Ann. Entomol. Soc. Am., 65: 1023-1025.
- IMAI H. T., 1974. — *B-chromosomes in the myrmicine ant, Leptothorax spinosus.* Chromosoma, 45: 431-444.