



Fig. 14 and 15. — *Pogonomyrmex* karyotypes: 14, *P. subnitidus* ♀. 15, *P. huachucanus* ♀. Scale line = 5  $\mu$ m.

Fig. 14 et 15. — Caryotypes de *Pogonomyrmex*: 14, *P. subnitidus* ♀. 15, *P. huachucanus* ♀. Echelle = 5  $\mu$ m.

- CROZIER R.H., 1977. — Genetic differentiation between populations of the ant *Aphaenogaster 'rudis'* in the southeastern United States. *Genetica*, 47 (1), 17-36.
- HAUSCHTECK E., 1961. — Die Chromosomen von fünf Ameisenarten. *Rev. Suisse Zool.*, 68, 218-223.
- IMAI H.T., 1966. — The chromosome observation techniques of ants and the chromosomes of Formicidae and Myrmecinae. *Acta Hymen.*, 2 (3), 119-131.
- IMAI H.T., KUBOTA M., 1972. — Karyological studies of Japanese ants (Hymenoptera: Formicidae). III. Karyotypes of nine species in Ponerinae, Formicinae, and Myrmecinae. *Chromosoma*, 37, 193-200.
- IMAI H.T., CROZIER R.H., TAYLOR R.W., 1977. — Karyotype evolution in Australian ants. *Chromosoma*, 59, 341-393.