

ACKNOWLEDGMENTS

Dr. Wojciech Pulawski (California Academy of Sciences) is gratefully acknowledged for loan of the *Willowsiella dispar* holotype, Kazuo Ogata and Philip S. Ward for comments, Colin Beaton and Helen Geier for scanning electron microscopy. Types of the new species were furnished by Harold Heatwole and Alan Andersen, for whom they are named. The author is an Honorary Research Fellow of CSIRO Division of Entomology.

SUMMARY

Romblonella heatwolei (Wyer Island, Torres Strait, Queensland) and *Willowsiella anderseni* (King Edward River, Western Australia) n. spp. are the first species of their genera to be reported from Australia. The genera are assigned to tribe Leptothoracini.

REFERENCES CITED

- BARONI-URBANI, C.
 1978. Materiali per una revisione dei *Leptothorax* neotropicale appartamenti al.sottogenere *Macromischa* Roger. Entomologica basil. 3: 395-618.
- BOLTON, B.
 1976. The ant tribe Tetramoriini (Hymenoptera: Formicidae). Constituent genera, review of small genera and revision of *Triglyphothrix* Forel. Bull. Br. Mus. Nat. Hist. (Ent). 34: 283-379.
 1981. A review of six minor genera of Myrmicinae in the Ethiopian zoogeographical region. Bull. Br. Mus. Nat. Hist. (Ent). 43: 245-307.
 1988. A review of *Paratopula* Wheeler, a forgotten genus of myrmicine ants (Hym., Formicidae). Entomologist's Mon. Mag. 124: 125-144.
 1982. Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotarsus*, *Messor* and *Cataulacus*. Bull. Br. Mus. Nat. Hist. (Ent). 45: 307-370.
- BROWN, W. L. JR.
 1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. In *Tropical forest ecosystems in Africa and South America: a comparative review*. Ed. Meggers, B. J., Ayensu, E. S. and Duckworth, W. D. Washington: Smithsonian Institution Press, pp. 161-185.
- EMERY, C.
 1897. Formicidarum species novae vel minus cognite in collectionae Musaei Nationalis Hungarici, quas in Nova-Guinea, Colonia Germanica, collectit L. Biró. Természetr. Fü. 20: 571-599.
 1922. Hymenoptera Fam. Formicidae Subfam. Myrmicinae in Wytzman, P. (ed.) Genera Insect. fasc. 174C; 192 pp (207-397).