



Figs. 1-8. SEM pictures of *Cerapachys sauteri* Forel.—1: Head, female. 2: Mandible, female. 3: Trunk, worker. 4: Petiole and postpetiole, worker. 5: Gaster, worker. 6: Surface structure of first gastric tergite, worker. 7: Trunk and petiole, male. 8: Terminal part of gaster, male, ventral view (SG, subgenital plate; PM, paramere; AD, aedeagus).

petiole and postpetiole strongly costate longitudinally; gaster subopaque and moderately punctate. Antennae, mandibles and legs opaque and weakly shagreened. Suberect hairs abundant, longest hair of trunk about 0.23 mm in length.

Color black; mandibles, antennae and legs blackish brown to brown.

*Description of male.* HL 1.10–1.15 mm; HW

1.23–1.25 mm; SL 0.40 mm; CI 109–111; SI 32–33; WL 2.30–2.45 mm; DTW 1.20–1.40 mm; PL 0.85 mm; PH 0.75–0.90 mm; DPW 0.75–0.78 mm; TL 6.5–7.0 mm. (Three individuals were measured.)

Head, excluding eyes, narrower than trunk; occipital border convex in full face view. Mandibles subtriangular, with 7 or 8 minute indistinct denticles. Anterior border of clypeus straight.