

Cerapachys biroi Forel, 1907

- Cerapachys biroi* Forel, 1907, p. 7.
Cerapachys silvestrii Wheeler, 1909, p. 269.
Cerapachys sinensis Wheeler, 1928, p. 3.
Cerapachys seini Mann, 1931, p. 440.
Cerapachys ierensis Weber, 1939, p. 94.

Material examined. Six workers, Chipen, Taitung City, 21-VII-1282, M. Terayama leg.

This is the first record of this species from Taiwan.

Distribution. Widely distributed in the Oriental region, and also introduced to Pacific islands and West Indies.

Cerapachys longitarsus (Mayr, 1878)

- Lioponera longitarsus* Mayr, 1878, p. 667.
Lioponera longitarsus var. *australis* Forel, 1895, p. 422.
Lioponera parva Forel, 1900, p. 330.
Lioponera bicolor Wheeler & Chapman, 1925, p. 54.
Phyracaces pygmaeus Clark, 1934, p. 26.
Cerapachys longitarsus: Brown, 1973, p. 181; 1975, p. 23.

No additional material was available.

Locarities previously recorded. Henchun, Pin-tung Hsien (=Kankau); Anping, Tainan City (=Anpin).

Distribution. India to the Philippines, North Australia.

Cerapachys reticulatus Emery, 1923

- Cerapachys reticulatus* Emery, 1923, p. 60.
No additional material was available. This species is known only by winged queen.

Locality previously recorded. Zhuqi Xiang, Chiai Hsien (=Suisharyo).

Distribution. Taiwan.

References

- Brown, W. L., 1973. A comparison of the hylaean and Congo-West Africa rain forest and faunas. In Tropical forest ecosystems in Africa and South America: A comparative review (B. J. Meggers, E. S. Ayensu, and W. D. Duckworth, eds.), pp. 161-185. Smiths. Inst. Press, Wash., D. C.
— 1975. Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmeceini, Acanthostichini and Aenictogitini. Search, 5: 1-115.
Clark, J., 1934. New Australian ants. Mem. nat. Mus. Vict., 8: 21-47.

- Emery, C., 1923. Einige exotische Ameisen des Deutschen Entomologischen Institutes. Ent. Mitt., 12: 60-62.
Forel, A., 1895. Nouvelles fourmis de l'Imerina Oriental (Moramanga, etc.). Ann. Soc. Ent. Belg., 39: 243-251. (Indirectly cited.)
— 1900. Les formicides de l'Empire des Indes ed de Ceylan. Part VII. J. Bombay Nat. Hist. Soc., 13: 303-332.
— 1907. Formicides du Musée National Hongrois. Ann. Mus. Nat. Hung., 5: 1-42.
— 1913. H. Sauter's Formosa-Ausbeute: Formicidae II. Arch. Naturg., (A) 79: 183-202.
Mann, W. M., 1931. A new ant from Porto Rico. J. Wash. Acad. Sci., 21: 440-441.
Mayr, G., 1878. Beiträge zur Ameisen-Fauna Asiens. Verh. Zool.-bot. Ges. Wien, 28: 645-686.
Weber, N. A., 1939. New ants of rare genera and a new genus of ponerine ants. Ann. Ent. Soc. Amer., 32: 91-104.
Wheeler, W. M., 1909. Ants collected by Prof. F. Silvestri in the Hawaiian Islands. Boll. Lab. Zool. Gen. Agr. Portici, 3: 269-272.
— 1928. Ants collected by Professor F. Silvestri in China. Boll. Lab. Zool. Gen. Agr. Portici, 22: 3-38.
Wheeler, W. M., and Chapman, J. W., 1925. The ants of the Philippine Islands. Part I. Dorylinae and Ponerinae. Philip. J. Sci., 28: 49-73.

(Terayama, M.: Toho Educational Institution. Chofu, Tokyo, 182 Japan. Kubota, S.: Akagidai Senior High School. Shinjuku-ku, Tokyo, 162 Japan. Sakai, H.: Oh! Elementary School. Ashigara-kami-gun, Kanagawa, 258 Japan. Kawazoe, A.: Yowa Junior High School. Kuwana, Mie, 511 Japan.)

Cerapachys sauteri Forel の再発見および台湾産 *Cerapachys* 属について

寺山 守・久保田敏・酒井春彦・川添昭夫

Cerapachys sauteri Forel, 1913 は台湾から職蟻によつて記載されたアリであるが、記載以降、本種の報告はなく、また模式標本も所在不明の状態であった。今回新たに本種の職蟻とともに雌、雄を検査することができたので、これらの記載を行つた。また本種が Brown (1973) の *crawleyi* グループに属することを確認するとともに、本グループの分布図を示した。さらに *C. biroi* Forel を台湾から初めて記録するとともに、台湾産 *Cerapachys* 属の種の検索表を提供した。

(寺山 守: 182 調布市若葉町 1-41-1 桐朋学園. 久保田 敏: 162 新宿区山吹町 81 赤城台高校. 酒井春彦: 258 神奈川県足柄上郡大井町金子 1436 大井小学校. 川添昭夫: 511 桑名市江場 7 陽和中学校)