



Figs. 1–8. *Camponotus yamaokai* sp. nov. (minor worker and major worker). — 1, Minor worker, head, frontal view; 2, minor worker, lateral view; 3, major worker, head, frontal view; 4, major worker, lateral view; 5, minor worker, mandible; 6, minor worker, petiole, lateral view (PSH: petiolar scale height, PSL: petiolar scale length); 7, minor worker, petiolar scale, dorsal view (DSW: dorsal petiolar scale width); 8, major worker petiolar scale, dorsal view.

two yellow spots.

*Major worker.* HL 1.28–1.38 mm; HW 1.28–1.38 mm; SL 0.88–0.95 mm; CI 96–106; SI 67–73; WL 1.55–1.70 mm; AW 0.83–0.90 mm; AI 53–58; PSL 0.16–0.18 mm; PSH 0.38–0.55 mm; DSW 0.49–0.55 mm; SCI 37–47; SWI 56–61; TL 4.8–5.3 mm. (Twenty individuals were measured.)

Head (Fig. 3) rectangular, with almost parallel sides and straight occipital border in front view. Mandibles strong, with rounded outer margin and five distinct teeth. Anterior border of clypeus truncated, but weakly convex. Eyes (Fig. 15) convex, 0.25–0.28 mm in maximum diameter. Antennal scapes slightly exceeding the occipital border of head.

Alitrunk (Fig. 4) massive, with weakly convex pro- and mesonotal dorsa and